Subject Index

Acid Neutralization
Acid neutralization of precipitation in Northern China, 204

Activated Carbon
Effects of activated carbon surface properties on the adsorption of volatile organic compounds, 1196
Preparation, characterization, and phenol adsorption of activated carbons from oxytetracycline bacterial residue, 1394

Activated Carbon Adsorbent
Breakthrough indicator for aromatic VOCs using needle trap samplers for activated carbon adsorbent, 946

Activated Carbon Particles
Toluene vapor capture by activated carbon particles in a dual gas-solid cyclone system, 924

Adsorbents
Analysis and comparison of inertinite-derived adsorbent with conventional adsorbents, 489

Adsorption
Adsorption Behavior of Toluene on Modified 1X Molecular Sieves, 1227
Effects of activated carbon surface properties on the adsorption of volatile organic compounds, 1196
Experimental study on removals of SO_2 and NO_x using adsorption of activated carbon/microwave desorption, 1012

Aeration
The emission patterns of volatile organic compounds during aerobic biotreatment of municipal solid waste using continuous and intermittent aeration, 461

Aerobic Biotreatment
The emission patterns of volatile organic compounds during aerobic biotreatment of municipal solid waste using continuous and intermittent aeration, 461

Aerosol
An inexpensive active optical remote sensing instrument for assessing aerosol distributions, 198
Prediction of daily fine particulate matter concentrations using aerosol optical depth retrievals from the Geostationary Operational Environmental Satellite (GOES), 1022

Aging
Aging of secondary organic aerosol from α-pinene ozonolysis: Roles of hydroxyl and nitrate radicals, 1359

Agricultural Area
Potential assessment of establishing a renewable energy plant in a rural agricultural area, 662

Air
Air impacts from three alternatives for producing JP-8 jet fuel, 1182
Application of ultraviolet light-emitting diode photocatalysis to remove volatile organic compounds from indoor air, 1032
Energy supplies and future engines for land, sea, and air, 607, 1233
Measurement of BTEX (benzene, toluene, ethybenzene, and xylene) levels at urban and semirural areas of Algiers City using passive air samplers, 1370
Near-source air quality impacts of large olefin flares, 978
Speciated arsenic in air: Measurement methodology and risk assessment considerations, 2

Airborne Ultrafine Particles
Assessment of exposure to airborne ultrafine particles in the urban environment of Lisbon, Portugal, 373
Exposure to airborne ultrafine particles from cooking in Portuguese homes, 1116

Aircraft Auxiliary Power Unit (APU)
Determination of the emissions from an aircraft auxiliary power unit (APU) during the Alternative Aviation Fuel Experiment (AAFEX), 420

Air Diffusion System
Effect of biocover equipped with a novel passive air diffusion system on microbial methane oxidation and community of methanotrophs, 278

Air Pollutants
Modeling an air pollution episode in northwestern United States: Identifying the effect of nitrogen oxide and volatile organic compound emission changes on air pollutants formation using direct sensitivity analysis, 1150

Air Pollution
Ambient air pollution and term birth weight in Texas from 1998 to 2004, 1285
An examination of boundary layer structure under the influence of the gap winds in Urumqi, China, during air pollution episode in winter, 26
Modeling an air pollution episode in northwestern United States: Identifying the effect of nitrogen oxide and volatile organic compound emission changes on air pollutants formation using direct sensitivity analysis, 1150
A reanalysis of fine particulate matter air pollution *versus* life expectancy in the United States, 989

Air Pollution Emissions Controls
Resolving the interactions between population density and air pollution emissions controls in the San Joaquin Valley, USA, 566

Air Quality
Development of risk-based air quality management strategies under impacts of climate change, 557
Review of Singapore's air quality and greenhouse gas emissions: Current situation and opportunities, 625

Air Quality Regulations
Historical trends in the mass and chemical species concentrations of coarse particulate matter in the Los Angeles Basin and relation to sources and air quality regulations, 541

Air Quality Zones
Determination of air quality zones in Turkey, 408

Alternative Aviation Fuel Experiment (AAFEX)
Determination of the emissions from an aircraft auxiliary power unit (APU) during the Alternative Aviation Fuel Experiment (AAFEX), 420

Ambient Air
Polycyclic aromatic hydrocarbon burden in ambient air in selected Niger Delta communities in Nigeria, 18

Ambient Air Pollution
Ambient air pollution and term birth weight in Texas from 1998 to 2004, 1285

Ammonia
Development of regenerative dye impregnated mesoporous silica materials for assessing exposure to ammonia, 838

Ammonia Emissions
Reducing ammonia emissions from laying-hen houses through dietary manipulation, 160

Animal Buildings
Comparison of six particle size distribution models on the goodness-of-fit to particulate matter sampled from animal buildings, 725

Animal Feed
Hydrothermal treatment for inactivating some hygienic microbial indicators from food waste–amended animal feed, 810

Aromatic Hydrocarbon
Polycyclic aromatic hydrocarbon burden in ambient air in selected Niger Delta communities in Nigeria, 18

Aromatic VOC
Breakthrough indicator for aromatic VOCs using needle trap samplers for activated carbon adsorbent, 946

Arsenic
Speciated arsenic in air: Measurement methodology and risk assessment considerations, 2

Artificial Lightweight Aggregates
Resource recycling through artificial lightweight aggregates from sewage sludge and derived ash using boric acid flux to lower co-melting temperature, 262

Ash
Resource recycling through artificial lightweight aggregates from sewage sludge and derived ash using boric acid flux to lower co-melting temperature, 262

Atmospheric Dispersion Modeling System
Simulating industrial emissions using Atmospheric Dispersion Modeling System: Model performance and source emission factors, 336

Atmospheric Fine Particles
A comparative study on the varying exposure to atmospheric fine and coarse particles under urban and rural conditions, 1313

Atmospheric Mercury Emissions
Atmospheric mercury emissions from waste combustions measured by continuous monitoring devices, 686

Control strategies of atmospheric mercury emissions from coal-fired power plants in China, 576

Attapulgite
The addition of modified attapulgite reduces the emission of nitrous oxide and ammonia from aerobically composted chicken manure, 1174

B-0, 10, 20
Comparison of on-road emissions for B-0, B-10, and B-20 in transit buses, 443

***Bacillus Anthracis* Surrogate Spores**
Thermal inactivation of *Bacillus anthracis* surrogate spores in a bench-scale enclosed landfill gas flare, 151

Bacterial Residue
Characteristics of penicillin bacterial residue, 485

Baghouse Filter
The evolution and impact of testing baghouse filter performance, 916

Beef Cattle Feedlots
Particulate matter emission rates from beef cattle feedlots in Kansas—Reverse dispersion modeling, 350

Bench-Scale Investigation
Mitigation of odor causing emissions—Bench-scale investigation, 1423

Bioaerosol Samplers
A wind tunnel test of newly developed personal bioaerosol samplers, 828

Biochar
Effectiveness and mechanisms of hydrogen sulfide adsorption by camphor-derived biochar, 873

Bioethanol-Gasoline Fuel Blends
Bioethanol–gasoline fuel blends: Exhaust emissions and morphological characterization of particulate from a moped engine, 888

Biofilm Reactor
Optimization of a horizontal-flow biofilm reactor for the removal of methane at low temperatures, 1166

Biofilter
Full-scale biofilter reduction efficiencies assessed using portable 24-hour sampling units, 170

Optimizing nutrient supply in a rotatory-switching biofilter for toluene vapor treatment, 451

Transformation of microflora during degradation of gaseous toluene in a biofilter detected using PCR-DGGE, 748

Biogas Generation
Effect of biogas generation on radon emissions from landfills receiving radium-bearing waste from shale gas development, 1040

Biogas Slurry Liquid
Effects of foliar application with compost tea and filtrate biogas slurry liquid on yield and fruit quality of Washington navel orange *(Citrus sinenesis Osbeck)* trees, 767

Biotrickling Filter
Removal of nitric oxide in a biotrickling filter under thermophilic condition using *Chelatococcus daeguensis*, 509

Black Carbon
Multiple-year black carbon measurements and source apportionment using Delta-C in Rochester, New York, 880

Black Carbon Particulate Matter Emission Factors
Black carbon particulate matter emission factors for buoyancy-driven associated gas flares, 307

Boric Acid Flux
Resource recycling through artificial lightweight aggregates from sewage sludge and derived ash using boric acid flux to lower co-melting temperature, 262

Boundary Layer Structure
An examination of boundary layer structure under the influence of the gap winds in Urumqi, China, during air pollution episode in winter, 26

BTEX
Measurement of BTEX (benzene, toluene, ethybenzene, and xylene) levels at urban and semirural areas of Algiers City using passive air samplers, 1370

Building Materials
Determination of partition and diffusion coefficients of formaldehyde in selected building materials and impact of relative humidity, 671

Buoyancy-Driven Associated Gas Flares
Black carbon particulate matter emission factors for buoyancy-driven associated gas flares, 307

Camphor-Derived Biochar
Effectiveness and mechanisms of hydrogen sulfide adsorption by camphor-derived biochar, 873

Canadian Agriculture
Trends in primary particulate matter emissions from Canadian agriculture, 737

Cancer Values of Prevented Fatalities
Cancer values of prevented fatalities (VPFs), one size does not fit all: The benefits of contaminated site cleanups in Italy, 783

Carbon/Microwave Desorption
Experimental study on removals of SO_2 and NO_x using adsorption of activated carbon/microwave desorption, 1012

Carbonyl Emissions
Carbonyl emissions from vehicular exhausts sources in Hong Kong, 221

Catalytic Reduction
The effects of manganese precursors on Mn-based/TiO_2 catalysts for catalytic reduction of NO with NH_3, 271

Cement Clinkers
Effects of sintering atmosphere on cement clinkers produced from chromium-bearing sludge, 587

Chars
Physicochemical properties of chars at different treatment temperatures, 235

Chelatococcus Daeguensis
Removal of nitric oxide in a biotrickling filter under thermophilic condition using *Chelatococcus daeguensis*, 509

Chicken Manure
The addition of modified attapulgite reduces the emission of nitrous oxide and ammonia from aerobically composted chicken manure, 1174

Chromium-Bearing Sludge
Effects of sintering atmosphere on cement clinkers produced from chromium-bearing sludge, 587

Clean Development Mechanism Application
Benefits of clean development mechanism application on the life cycle assessment perspective: A case study in the palm oil industry, 299

Climate Change
Development of risk-based air quality management strategies under impacts of climate change, 557

Evaluating the effects of climate change on summertime ozone using a relative response factor approach for policymakers, 1061

Coal Combustion Flue Gas
Effect of solution pH on SO_2, NOx, and Hg removal from simulated coal combustion flue gas in an oxidant-enhanced wet scrubber, 212

Coal-Fired Power Plants
Control strategies of atmospheric mercury emissions from coal-fired power plants in China, 576

Coarse Particulate Matter
Historical trends in the mass and chemical species concentrations of coarse particulate matter in the Los Angeles Basin and relation to sources and air quality regulations, 541

Coke Oven Plant
Temporal and spatial variations of polycyclic aromatic hydrocarbon concentrations around a coke oven plant, 1003

Combustion Flue Gas
Gas-phase elemental mercury removal in a simulated combustion flue gas using TiO_2 with fluorescent light, 1208

Co-Melting Technology
Co-melting technology in resource recycling of sludge derived from stone processing, 1449

Co-Melting Temperature
Resource recycling through artificial lightweight aggregates from sewage sludge and derived ash using boric acid flux to lower co-melting temperature, 262

Community Functional Diversity
Comparison of the purification performance and microbial community functional diversity in flow directional- switching and unidirectional-flow biotrickling filters, 1203

Compost Tea
Effects of foliar application with compost tea and filtrate biogas slurry liquid on yield and fruit quality of Washington navel orange *(Citrus sinenesis Osbeck)* trees, 767

Consumer Products
Emissions of *p*-dichlorobenzene and naphthalene from consumer products, 1075

Contaminated Site Cleanups
Cancer values of prevented fatalities (VPFs), one size does not fit all: The benefits of contaminated site cleanups in Italy, 783

Contiguous United States
Source attribution for mercury deposition in the contiguous United States: Regional difference and seasonal variation, 52

Control Strategies
Control strategies of atmospheric mercury emissions from coal-fired power plants in China, 576

Ozone and its potential control strategy for Chon Buri city, Thailand, 1411

Cooking
Exposure to airborne ultrafine particles from cooking in Portuguese homes, 1116

Corrosive Gas
Integration of a nonmetallic electrostatic precipitator and a wet scrubber for improved removal of particles and corrosive gas cleaning in semiconductor manufacturing industries, 905

Cross-Linked Biopolymeric Glass Wastes
Feasibility study on cross-linked biopolymeric concrete encapsulating selenium glass wastes, 898

Dairy Operation
Comparison of seasonal phenol and p-cresol emissions from ground-level area sources in a dairy operation in central Texas, 381

Deep-Pit Swine Finishing
Emissions monitoring at a deep-pit swine finishing facility: Research methods and system performance, 1264

Degradation
Transformation of microflora during degradation of gaseous toluene in a biofilter detected using PCR-DGGE, 748

Delta-C
Multiple-year black carbon measurements and source apportionment using Delta-C in Rochester, New York, 880

Dietary Manipulation
Reducing ammonia emissions from laying-hen houses through dietary manipulation, 160

Diffusion Coefficients
Determination of partition and diffusion coefficients of formaldehyde in selected building materials and impact of relative humidity, 671

Dilution of Vehicle Exhaust Pollutants
Experimental studies of the dilution of vehicle exhaust pollutants by environment-protecting pervious pavement, 92

Dioxins
Dioxins and their fingerprint in size-classified fly ash fractions from municipal solid waste incinerators in China—Mechanical grate and fluidized bed units, 717

Dispersion Modeling
Estimation of hydrogen sulfide emission rates at several wastewater treatment plants through experimental concentration measurements and dispersion modeling, 758

NO_2 and SO_2 dispersion modeling and relative roles of emission sources over Map Ta Phut industrial area, Thailand, 932

Diurnal Pattern of PM_{10}
Relative roles of emissions and meteorology in the diurnal pattern of urban PM_{10}: Analysis of the daylight saving time effect, 642

Domestic Emission
Sensitivity and linearity analysis of ozone in East Asia: The effects of domestic emission and intercontinental transport, 1102

Driver Exposure
Driver exposure to particulate matter in Bangkok, 64

Dual Gas-Solid Cyclone System
Toluene vapor capture by activated carbon particles in a dual gas-solid cyclone system, 924

Dust Levels
Application of neutral electrolyzed water spray for reducing dust levels in a layer breeding house, 1329

Dye
Development of regenerative dye impregnated mesoporous silica materials for assessing exposure to ammonia, 838

EMFAC2007
Comparison of the MOVES2010a, MOBILE6.2, and EMFAC2007 mobile source emission models with on-road traffic tunnel and remote sensing measurements, 1134

Emissions
Determination of the emissions from an aircraft auxiliary power unit (APU) during the Alternative Aviation Fuel Experiment (AAFEX), 420

Emission Contributions
A new approach to characterize emission contributions from area sources during optical remote sensing technique testing, 1403

Emission Factors
A tunnel study to estimate emission factors from mobile sources in Monterrey, Mexico, 1431

Emission Factor Uncertainty
Quantification of emission factor uncertainty, 287

Emission Patterns
The emission patterns of volatile organic compounds during aerobic biotreatment of municipal solid waste using continuous and intermittent aeration, 461

Emissions Monitoring
Emissions monitoring at a deep-pit swine finishing facility: Research methods and system performance, 1264

Emission Sources
NO_2 and SO_2 dispersion modeling and relative roles of emission sources over Map Ta Phut industrial area, Thailand, 932

Energy Source
Evaluation of a sustainable remediation option: Beneficial reuse of petroleum-contaminated sediment as an energy source, 1277

Energy Supplies
Energy supplies and future engines for land, sea, and air, 607, 1233

Environmental Impact
An economic evaluation and assessment of environmental impact of the municipal solid waste management system for Taichung City in Taiwan, 527

Environmental Management Problem
A generalized fuzzy linear programming approach for environmental management problem under uncertainty, 72

Environment-Protecting Pavement
Experimental studies of the dilution of vehicle exhaust pollutants by environment-protecting pervious pavement, 92

Exhaust Emissions
Bioethanol–gasoline fuel blends: Exhaust emissions and morphological characterization of particulate from a moped engine, 888

Study of Miller timing on exhaust emissions of a hydrotreated vegetable oil (HVO)-fueled diesel engine, 1305

Exposure to Airborne Ultrafine Particles
Exposure to airborne ultrafine particles from cooking in Portuguese homes, 1116

Extreme Value Modeling
Extreme value modeling for the analysis and prediction of time series of extreme tropospheric ozone levels: A case study, 651

Fabric Filter
Ultrafine particle emission from incinerators: The role of the fabric filter, 103

Filter Cleaning
Influence of pleat geometry on filter cleaning in PTFE/glass composite filter, 1257

Fine Particulate Matter
Assessment of heterogeneity of metal composition of fine particulate matter collected from eight U.S. counties using principal component analysis, 773

Fine particulate matter and visibility in the Lake Tahoe Basin: Chemical characterization, trends, and source apportionment, 953

A novel methodology for determining low-cost fine particulate matter street sweeping routes, 242

Prediction of daily fine particulate matter concentrations using aerosol optical depth retrievals from the Geostationary Operational Environmental Satellite (GOES), 1022

A reanalysis of fine particulate matter air pollution *versus* life expectancy in the United States, 989

Fine Particulate Pollutants
Multiple comparisons of organic, microbial, and fine particulate pollutants in typical indoor environments: Diurnal and seasonal variation, 1380

Flowdirectional Switching
Comparison of the purification performance and microbial community functional diversity in flowdirectional- switching and unidirectional-flow biotrickling filters, 1203

Flue Gas Desulfurization
Mercury emission and plant uptake of trace elements during early stage of soil amendment using flue gas desulfurization materials, 139

Fluidized Bed Units
Dioxins and their fingerprint in size-classified fly ash fractions from municipal solid waste incinerators in China—Mechanical grate and fluidized bed units, 717

Fluorescent Light
Gas-phase elemental mercury removal in a simulated combustion flue gas using TiO_2 with fluorescent light, 1208

Fly Ash
Effect of thermal processing on the characteristics of incineration fly ash, 500

Fly Ash Fractions
Dioxins and their fingerprint in size-classified fly ash fractions from municipal solid waste incinerators in China—Mechanical grate and fluidized bed units, 717

Foliar Application
Effects of foliar application with compost tea and filtrate biogas slurry liquid on yield and fruit quality of Washington navel orange *(Citrus sinenesis Osbeck)* trees, 767

Food Waste
Hydrothermal treatment for inactivating some hygienic microbial indicators from food waste–amended animal feed, 810

Formaldehyde
Determination of partition and diffusion coefficients of formaldehyde in selected building materials and impact of relative humidity, 671

Fruit Quality
Effects of foliar application with compost tea and filtrate biogas slurry liquid on yield and fruit quality of Washington navel orange *(Citrus sinenesis Osbeck)* trees, 767

Fuel
Quantification of personal exposure concentrations to gasoline vehicle emissions in high-end exposure microenvironments: Effects of fuel and season, 1346

Fuel Switching
Demonstration of fuel switching on oceangoing vessels in the Gulf of Mexico, 1093

Full-Scale Biofilter Reduction
Full-scale biofilter reduction efficiencies assessed using portable 24-hour sampling units, 170

Future Engines
Energy supplies and future engines for land, sea, and air, 1233

Fuzzy Linear Programming
A generalized fuzzy linear programming approach for environmental management problem under uncertainty, 72

Gap Winds
An examination of boundary layer structure under the influence of the gap winds in Urumqi, China, during air pollution episode in winter, 26

Gas
Compositions and greenhouse gas emission factors of flared and vented gas in the Western Canadian Sedimentary Basin, 992

Gas Cleaning System
The importance of the location of sodium chlorite application in a multipollutant flue gas cleaning system, 707

Gaseous Toluene
Transformation of microflora during degradation of gaseous toluene in a biofilter detected using PCR-DGGE, 748

Gas Flares
Black carbon particulate matter emission factors for buoyancy-driven associated gas flares, 307

Gasoline Vehicle Emissions
Quantification of personal exposure concentrations to gasoline vehicle emissions in high-end exposure microenvironments: Effects of fuel and season, 1346

Geostationary Operational Environmental Satellite (GOES)
Prediction of daily fine particulate matter concentrations using aerosol optical depth retrievals from the Geostationary Operational Environmental Satellite (GOES), 1022

Greenhouse Gas Emissions
Compositions and greenhouse gas emission factors of flared and vented gas in the Western Canadian Sedimentary Basin, 992

Review of Singapore's air quality and greenhouse gas emissions: Current situation and opportunities, 625

Ground-Level Area Sources
Comparison of seasonal phenol and p-cresol emissions from ground-level area sources in a dairy operation in central Texas, 381

Heterogeneity
Assessment of a regulatory model's performance relative to large spatial heterogeneity in observed ozone in Houston, Texas, 696

High-End Exposure Microenvironments
Quantification of personal exposure concentrations to gasoline vehicle emissions in high-end exposure microenvironments: Effects of fuel and season, 1346

8-Hour Ozone
Efficacy of recent state implementation plans for 8-hour ozone, 252

Hydrogen Sulfide
Characterization of the olfactory impact around a wastewater treatment plant: Optimization and validation of a hydrogen sulfide determination procedure based on passive diffusion sampling, 863

Hydrogen Sulfide Adsorption
Effectiveness and mechanisms of hydrogen sulfide adsorption by camphor-derived biochar, 873

Hydrogen Sulfide Emission
Estimation of hydrogen sulfide emission rates at several wastewater treatment plants through experimental concentration measurements and dispersion modeling, 758

Hydrogen Sulfide Removal
Middle-thermophilic sulfur-oxidizing bacteria *Thiomonas* sp. RAN5 strain for hydrogen sulfide removal, 38

Hydrothermal Treatment
Hydrothermal treatment for inactivating some hygienic microbial indicators from food waste–amended animal feed, 810

Hydrotreated Vegetable Oil (HVO)-Fueled Diesel Engine
Study of Miller timing on exhaust emissions of a hydrotreated vegetable oil (HVO)-fueled diesel engine, 1305

Hydroxyl Radicals
Aging of secondary organic aerosol from α-pinene ozonolysis: Roles of hydroxyl and nitrate radicals, 1359

Incineration Fly Ash
Effect of thermal processing on the characteristics of incineration fly ash, 500

Incinerators
Ultrafine particle emission from incinerators: The role of the fabric filter, 103

Indoor Air Pollutants Emitted
A real-time monitoring and assessment method for calculation of total amounts of indoor air pollutants emitted in subway stations, 517

Indoor Environments
Multiple comparisons of organic, microbial, and fine particulate pollutants in typical indoor environments: Diurnal and seasonal variation, 1380

Industrial Emissions
Simulating industrial emissions using Atmospheric Dispersion Modeling System: Model performance and source emission factors, 336

Inertinite-Derived Adsorbent
Analysis and comparison of inertinite-derived adsorbent with conventional adsorbents, 489

Intercontinental Transport
Sensitivity and linearity analysis of ozone in East Asia: The effects of domestic emission and intercontinental transport, 1102

Iron Degradation
Microwave-induced nanoscale zero-valent iron degradation of perchloroethylene and pentachlorophenol, 1443

JP-8 Jet Fuel
Air impacts from three alternatives for producing JP-8 jet fuel, 1182

Lake Tahoe Basin
Fine particulate matter and visibility in the Lake Tahoe Basin: Chemical characterization, trends, and source apportionment, 953

Land
Energy supplies and future engines for land, sea, and air, 607, 1233

Landfills
Effect of biogas generation on radon emissions from landfills receiving radium-bearing waste from shale gas development, 1040

Methane emissions from 20 landfills across the United States using vertical radial plume mapping, 183

Landfill Gas Flare
Thermal inactivation of *Bacillus anthracis* surrogate spores in a bench-scale enclosed landfill gas flare, 151

Lattice Oxygen
Effect of the Mn oxidation state and lattice oxygen in Mn-based TiO_2 catalysts on the low-temperature selective catalytic reduction of NO by NH_3, 1085

Layer Breeding House
Application of neutral electrolyzed water spray for reducing dust levels in a layer breeding house, 1329

Layer House
Field evaluation of particulate matter measurements using tapered element oscillating microbalance in a layer house, 322

Laying-Hen Houses
Reducing ammonia emissions from laying-hen houses through dietary manipulation, 160

Life Cycle Assessment Perspective
Benefits of clean development mechanism application on the life cycle assessment perspective: A case study in the palm oil industry, 299

Life Expectancy
A reanalysis of fine particulate matter air pollution *versus* life expectancy in the United States, 989

Light-Emitting Diode Photocatalysis
Application of ultraviolet light-emitting diode photocatalysis to remove volatile organic compounds from indoor air, 1032

Los Angeles Basin
Historical trends in the mass and chemical species concentrations of coarse particulate matter in the Los Angeles Basin and relation to sources and air quality regulations, 541

Management Strategies
Development of risk-based air quality management strategies under impacts of climate change, 557

Manganese Precursors
The effects of manganese precursors on Mn-based/TiO_2 catalysts for catalytic reduction of NO with NH_3, 271

Mechanical Grate Units
Dioxins and their fingerprint in size-classified fly ash fractions from municipal solid waste incinerators in China—Mechanical grate and fluidized bed units, 717

Mercuric Chloride Adsorption
Enhanced mercuric chloride adsorption onto sulfur-modified activated carbons derived from waste tires, 799

Mercury
Effect of solution pH on SO_2, NOx, and Hg removal from simulated coal combustion flue gas in an oxidant-enhanced wet scrubber, 212

Gas-phase elemental mercury removal in a simulated combustion flue gas using TiO_2 with fluorescent light, 1208

Mercury Deposition
Source attribution for mercury deposition in the contiguous United States: Regional difference and seasonal variation, 52

Mercury Emissions
Atmospheric mercury emissions from waste combustions measured by continuous monitoring devices, 686

Mercury emission and plant uptake of trace elements during early stage of soil amendment using flue gas desulfurization materials, 139

Mesoporous Silica Materials
Development of regenerative dye impregnated mesoporous silica materials for assessing exposure to ammonia, 838

Metal Composition, Heterogeneity
Assessment of heterogeneity of metal composition of fine particulate matter collected from eight U.S. counties using principal component analysis, 773

Methane, Removal
Optimization of a horizontal-flow biofilm reactor for the removal of methane at low temperatures, 1166

Methane Emissions
Methane emissions from 20 landfills across the United States using vertical radial plume mapping, 183

Methanotrophs
Effect of biocover equipped with a novel passive air diffusion system on microbial methane oxidation and community of methanotrophs, 278

Microbial Indicators
Hydrothermal treatment for inactivating some hygienic microbial indicators from food waste–amended animal feed, 810

Microbial Methane Oxidation
Effect of biocover equipped with a novel passive air diffusion system on microbial methane oxidation and community of methanotrophs, 278

Microbial Pollutants
Multiple comparisons of organic, microbial, and fine particulate pollutants in typical indoor environments: Diurnal and seasonal variation, 1380

Microflora
Transformation of microflora during degradation of gaseous toluene in a biofilter detected using PCR-DGGE, 748

Microwave-Induced Nanoscale Zero-Valent Iron Degradation
Microwave-induced nanoscale zero-valent iron degradation of perchloroethylene and pentachlorophenol, 1443

Middle-Thermophilic Sulfur-Oxidizing Bacteria
Middle-thermophilic sulfur-oxidizing bacteria *Thiomonas* sp. RAN5 strain for hydrogen sulfide removal, 38

Miller Timing
Study of Miller timing on exhaust emissions of a hydrotreated vegetable oil (HVO)-fueled diesel engine, 1305

Mn, Valence State
Improving the activity of Mn/TiO2 catalysts through control of the pH and valence state of Mn during their preparation, 362

Mn-Based/TiO_2 Catalysts
The effects of manganese precursors on Mn-based/TiO_2 catalysts for catalytic reduction of NO with NH_3, 271

Mn Oxidation State
Effect of the Mn oxidation state and lattice oxygen in Mn-based TiO_2 catalysts on the low-temperature selective catalytic reduction of NO by NH_3, 1085

Mn/TiO_2 Catalysts
Improving the activity of Mn/TiO_2 catalysts through control of the pH and valence state of Mn during their preparation, 362

MOBILE6.2
Comparison of the MOVES2010a, MOBILE6.2, and EMFAC2007 mobile source emission models with on-road traffic tunnel and remote sensing measurements, 1134

Mobile Monitoring System
A study on the characteristics of silt loading on paved roads in the Seoul metropolitan area using a mobile monitoring system, 846

Mobile Source Carbon Dioxide Emissions
A 2009 Mobile Source Carbon Dioxide Emissions Inventory for the University of Central Florida, 1050

Mobile Source Emission Models
Comparison of the MOVES2010a, MOBILE6.2, and EMFAC2007 mobile source emission models with on-road traffic tunnel and remote sensing measurements, 1134

Mobile Source Indicators
Development of outcome-based, multipollutant mobile source indicators, 431

Mobile Sources
A tunnel study to estimate emission factors from mobile sources in Monterrey, Mexico, 1431

Modeling
Assessment of a regulatory model's performance relative to large spatial heterogeneity in observed ozone in Houston, Texas, 696

Extreme value modeling for the analysis and prediction of time series of extreme tropospheric ozone levels: A case study, 651

Modeling an air pollution episode in northwestern United States: Identifying the effect of nitrogen oxide and volatile organic compound emission changes on air pollutants formation using direct sensitivity analysis, 1150

Particulate matter emission rates from beef cattle feedlots in Kansas—Reverse dispersion modeling, 350

Simulating industrial emissions using Atmospheric Dispersion Modeling System: Model performance and source emission factors, 336

Molecular Markers
Obtaining polycyclic aromatic hydrocarbon concentration ratios and molecular markers for residential wood combustion: Temuco, a case study, 44

Molecular Sieves
Adsorption Behavior of Toluene on Modified 1X Molecular Sieves, 1227

Moped Engine
Bioethanol–gasoline fuel blends: Exhaust emissions and morphological characterization of particulate from a moped engine, 888

MOVES2010a
Comparison of the MOVES2010a, MOBILE6.2, and EMFAC2007 mobile source emission models with on-road traffic tunnel and remote sensing measurements, 1134

Multipollutant
Near-road multipollutant profiles: Associations between volatile organic compounds and a tracer gas surrogate near a busy highway, 594

Multipollutant Flue Gas Cleaning System
The importance of the location of sodium chlorite application in a multipollutant flue gas cleaning system, 707

Multipollutant Mobile Source Indicators
Development of outcome-based, multipollutant mobile source indicators, 431

Multipollutant Profiles
Near-road multipollutant profiles: Associations between volatile organic compounds and a tracer gas surrogate near a busy highway, 594

Multispecies Remote Sensing
Multispecies remote sensing measurements of vehicle emissions on ShermanWay in Van Nuys, California, 1127

Municipal Solid Waste
The emission patterns of volatile organic compounds during aerobic biotreatment of municipal solid waste using continuous and intermittent aeration, 461

Municipal Solidwaste Disposal
Volatile organic compound emissions from municipal solidwaste disposal sites:Acase study of Mumbai, India, 398

Municipal Solid Waste Incinerators
Dioxins and their fingerprint in size-classified fly ash fractions from municipal solid waste incinerators in China—Mechanical grate and fluidized bed units, 717

Municipal Solid Waste Management System
An economic evaluation and assessment of environmental impact of the municipal solid waste management system for Taichung City in Taiwan, 527

Nanoparticles
Influence of particle location on the number of charges per charged nanoparticle at the outlet of a needle charger, 87

Naphthalene
Emissions of *p*-dichlorobenzene and naphthalene from consumer products, 1075

Near-Source Air Quality
Near-source air quality impacts of large olefin flares, 978

Near-Source Ozone
The potential near-source ozone impacts of upstream oil and gas industry emissions, 966

Needle Charger
Influence of particle location on the number of charges per charged nanoparticle at the outlet of a needle charger, 87

Needle Trap Samplers
Breakthrough indicator for aromatic VOCs using needle trap samplers for activated carbon adsorbent, 946

Negative Air Ion
Enhancement of the deposition of ultrafine secondary organic aerosols by the negative air ion and the effect of relative humidity, 1296

Neutral Electrolyzed Water Spray
Application of neutral electrolyzed water spray for reducing dust levels in a layer breeding house, 1329

NH_3
The addition of modified attapulgite reduces the emission of nitrous oxide and ammonia from aerobically composted chicken manure, 1174

Effect of the Mn oxidation state and lattice oxygen in Mn-based TiO_2 catalysts on the low-temperature selective catalytic reduction of NO by NH_3, 1085

The effects of manganese precursors on Mn-based/TiO_2 catalysts for catalytic reduction of NO with NH_3, 271

Niger Delta communities
Polycyclic aromatic hydrocarbon burden in ambient air in selected Niger Delta communities in Nigeria, 18

Nitrate Radicals
Aging of secondary organic aerosol from α-pinene ozonolysis: Roles of hydroxyl and nitrate radicals, 1359

NO
The addition of modified attapulgite reduces the emission of nitrous oxide and ammonia from aerobically composted chicken manure, 1174
Effect of the Mn oxidation state and lattice oxygen in Mn-based TiO_2 catalysts on the low-temperature selective catalytic reduction of NO by NH_3, 1085
The effects of manganese precursors on Mn-based/TiO_2 catalysts for catalytic reduction of NO with NH_3, 271
Modeling an air pollution episode in northwestern United States: Identifying the effect of nitrogen oxide and volatile organic compound emission changes on air pollutants formation using direct sensitivity analysis, 1150
Removal of nitric oxide in a biotrickling filter under thermophilic condition using *Chelatococcus daeguensis*, 509

NO_2
Assessing the spatial distribution of nitrogen dioxide in London, Ontario, 1335
NO_2 and SO_2 dispersion modeling and relative roles of emission sources over Map Ta Phut industrial area, Thailand, 932

Nonmetallic Electrostatic Precipitator
Integration of a nonmetallic electrostatic precipitator and a wet scrubber for improved removal of particles and corrosive gas cleaning in semiconductor manufacturing industries, 905

NOx
Effect of solution pH on SO_2, NOx, and Hg removal from simulated coal combustion flue gas in an oxidant-enhanced wet scrubber, 212
Experimental study on removals of SO_2 and NO_x using adsorption of activated carbon/microwave desorption, 1012

Nutrient Supply
Optimizing nutrient supply in a rotatory-switching biofilter for toluene vapor treatment, 451

Oceangoing Vessels
Demonstration of fuel switching on oceangoing vessels in the Gulf of Mexico, 1093

Odor Causing Emissions
Mitigation of odor causing emissions—Bench-scale investigation, 1423

OH Radicals
Rapidly eliminating pathogenic microorganisms in large air space using spraying •OH radicals, 393

Oil and Gas Industry Emissions
The potential near-source ozone impacts of upstream oil and gas industry emissions, 966

Olefin Flares
Near-source air quality impacts of large olefin flares, 978

Olfactory Impact
Characterization of the olfactory impact around a wastewater treatment plant: Optimization and validation of a hydrogen sulfide determination procedure based on passive diffusion sampling, 863

On-Road Emissions
Comparison of on-road emissions for B-0, B-10, and B-20 in transit buses, 443

On-Road Traffic Tunnel
Comparison of the MOVES2010a, MOBILE6.2, and EMFAC2007 mobile source emission models with on-road traffic tunnel and remote sensing measurements, 1134

On-Road Vehicle Emissions
Effects of improved spatial and temporal modeling of on-road vehicle emissions, 471

Optical Remote Sensing
An inexpensive active optical remote sensing instrument for assessing aerosol distributions, 198
A new approach to characterize emission contributions from area sources during optical remote sensing technique testing, 1403

Orange *(Citrus Sinenesis Osbeck)* Trees
Effects of foliar application with compost tea and filtrate biogas slurry liquid on yield and fruit quality of Washington navel orange *(Citrus sinenesis Osbeck)* trees, 767

Organic Aerosols
Aging of secondary organic aerosol from α-pinene ozonolysis: Roles of hydroxyl and nitrate radicals, 1359
Enhancement of the deposition of ultrafine secondary organic aerosols by the negative air ion and the effect of relative humidity, 1296

Organic Pollutants
Multiple comparisons of organic, microbial, and fine particulate pollutants in typical indoor environments: Diurnal and seasonal variation, 1380

Oscillating Microbalance
Field evaluation of particulate matter measurements using tapered element oscillating microbalance in a layer house, 322

Oxidant-Enhanced Wet Scrubber
Effect of solution pH on SO_2, NOx, and Hg removal from simulated coal combustion flue gas in an oxidant-enhanced wet scrubber, 212

Ozone
Assessment of a regulatory model's performance relative to large spatial heterogeneity in observed ozone in Houston, Texas, 696
Efficacy of recent state implementation plans for 8-hour ozone, 252
Evaluating the effects of climate change on summertime ozone using a relative response factor approach for policymakers, 1061
Ozone and its potential control strategy for Chon Buri city, Thailand, 1411
Sensitivity and linearity analysis of ozone in East Asia: The effects of domestic emission and intercontinental transport, 1102

Ozone Levels
Extreme value modeling for the analysis and prediction of time series of extreme tropospheric ozone levels: A case study, 651

Palm Oil Industry
Benefits of clean development mechanism application on the life cycle assessment perspective: A case study in the palm oil industry, 299

Particle Location
Influence of particle location on the number of charges per charged nanoparticle at the outlet of a needle charger, 87

Particle Size Distribution Models
Comparison of six particle size distribution models on the goodness-of-fit to particulate matter sampled from animal buildings, 725

Particulate Matter
Driver exposure to particulate matter in Bangkok, 64

Particulate Matter Emission Rates
Particulate matter emission rates from beef cattle feedlots in Kansas—Reverse dispersion modeling, 350

Particulate Matter Measurements
Field evaluation of particulate matter measurements using tapered element oscillating microbalance in a layer house, 322

Passive Diffusion Sampling
Characterization of the olfactory impact around a wastewater treatment plant: Optimization and validation of a hydrogen sulfide determination procedure based on passive diffusion sampling, 863

Pathogenic Microorganisms
Rapidly eliminating pathogenic microorganisms in large air space using spraying •OH radicals, 393

Paved Roads
A study on the characteristics of silt loading on paved roads in the Seoul metropolitan area using a mobile monitoring system, 846

PCR-DGGE
Transformation of microflora during degradation of gaseous toluene in a biofilter detected using PCR-DGGE, 748

***p*-Cresol Emissions**
Comparison of seasonal phenol and p-cresol emissions from ground-level area sources in a dairy operation in central Texas, 381

***p*-Dichlorobenzene**
Emissions of *p*-dichlorobenzene and naphthalene from consumer products, 1075

Penicillin Bacterial Residue
Characteristics of penicillin bacterial residue, 485

Pentachlorophenol
Microwave-induced nanoscale zero-valent iron degradation of perchloroethylene and pentachlorophenol, 1443

Perchloroethylene
Microwave-induced nanoscale zero-valent iron degradation of perchloroethylene and pentachlorophenol, 1443

Personal Exposure Concentrations
Quantification of personal exposure concentrations to gasoline vehicle emissions in high-end exposure microenvironments: Effects of fuel and season, 1346

Petroleum-Contaminated Sediment
Evaluation of a sustainable remediation option: Beneficial reuse of petroleum-contaminated sediment as an energy source, 1277

pH
Effect of solution pH on SO_2, NOx, and Hg removal from simulated coal combustion flue gas in an oxidant-enhanced wet scrubber, 212

Improving the activity of Mn/TiO2 catalysts through control of the pH and valence state of Mn during their preparation, 362

phenol
Comparison of seasonal phenol and p-cresol emissions from ground-level area sources in a dairy operation in central Texas, 381

Phenol Adsorption
Preparation, characterization, and phenol adsorption of activated carbons from oxytetracycline bacterial residue, 1394

Photocatalysis
Application of ultraviolet light-emitting diode photocatalysis to remove volatile organic compounds from indoor air, 1032

α-Pinene Ozonolysis
Aging of secondary organic aerosol from α-pinene ozonolysis: Roles of hydroxyl and nitrate radicals, 1359

Plant Uptake
Mercury emission and plant uptake of trace elements during early stage of soil amendment using flue gas desulfurization materials, 139

Pleat Geometry
Influence of pleat geometry on filter cleaning in PTFE/glass composite filter, 1257

Plumes
Evaluating the performance of the horizontal radial plume mapping technique for locating multiple plumes, 1249

Plume Mapping
Evaluating the performance of the horizontal radial plume mapping technique for locating multiple plumes, 1249

Methane emissions from 20 landfills across the United States using vertical radial plume mapping, 183

$PM_{2.5}$
Chemical characterization and factor analysis of $PM_{2.5}$ in two sites of Monterrey, Mexico, 817

$PM_{2.5}$ Chemical Compositions
Winter and Summer $PM_{2.5}$ Chemical Compositions in Fourteen Chinese Cities, 1214

PM_{10}
Relative roles of emissions and meteorology in the diurnal pattern of urban PM_{10}: Analysis of the daylight saving time effect, 642

Policymakers
Evaluating the effects of climate change on summertime ozone using a relative response factor approach for policymakers, 1061

Polycyclic Aromatic Hydrocarbon
Obtaining polycyclic aromatic hydrocarbon concentration ratios and molecular markers for residential wood combustion: Temuco, a case study, 44

Temporal and spatial variations of polycyclic aromatic hydrocarbon concentrations around a coke oven plant, 1003

Polycyclic aromatic hydrocarbon
Polycyclic aromatic hydrocarbon burden in ambient air in selected Niger Delta communities in Nigeria, 18

Population Density
Resolving the interactions between population density and air pollution emissions controls in the San Joaquin Valley, USA, 566

Precipitation
Acid neutralization of precipitation in Northern China, 204

Primary Particulate Matter Emissions
Trends in primary particulate matter emissions from Canadian agriculture, 737

Principal Component Analysis
Assessment of heterogeneity of metal composition of fine particulate matter collected from eight U.S. counties using principal component analysis, 773

PTFE/Glass Composite Filter
Influence of pleat geometry on filter cleaning in PTFE/glass composite filter, 1257

Quantification
Quantification of emission factor uncertainty, 287

Quantification of personal exposure concentrations to gasoline vehicle emissions in high-end exposure microenvironments: Effects of fuel and season, 1346

Radium-Bearing Waste
Effect of biogas generation on radon emissions from landfills receiving radium-bearing waste from shale gas development, 1040

RAN5 Strain
Middle-thermophilic sulfur-oxidizing bacteria *Thiomonas* sp. RAN5 strain for hydrogen sulfide removal, 38

Regenerative Dye
Development of regenerative dye impregnated mesoporous silica materials for assessing exposure to ammonia, 838

Regulatory Model's Performance
Assessment of a regulatory model's performance relative to large spatial heterogeneity in observed ozone in Houston, Texas, 696

Relative Humidity
Determination of partition and diffusion coefficients of formaldehyde in selected building materials and impact of relative humidity, 671

Enhancement of the deposition of ultrafine secondary organic aerosols by the negative air ion and the effect of relative humidity, 1296

Remediation
Evaluation of a sustainable remediation option: Beneficial reuse of petroleum-contaminated sediment as an energy source, 1277

Remote Sensing
An inexpensive active optical remote sensing instrument for assessing aerosol distributions, 198

Comparison of the MOVES2010a, MOBILE6.2, and EMFAC2007 mobile source emission models with on-road traffic tunnel and remote sensing measurements, 1134

Multispecies remote sensing measurements of vehicle emissions on ShermanWay in Van Nuys, California, 1127

A new approach to characterize emission contributions from area sources during optical remote sensing technique testing, 1403

Renewable Energy Plant
Potential assessment of establishing a renewable energy plant in a rural agricultural area, 662

Resource Recycling
Co-melting technology in resource recycling of sludge derived from stone processing, 1449

Resource recycling through artificial lightweight aggregates from sewage sludge and derived ash using boric acid flux to lower co-melting temperature, 262

Reverse Dispersion Modeling
Particulate matter emission rates from beef cattle feedlots in Kansas—Reverse dispersion modeling, 350

Risk Assessment
Speciated arsenic in air: Measurement methodology and risk assessment considerations, 2

Rotatory-Switching Biofilter
Optimizing nutrient supply in a rotatory-switching biofilter for toluene vapor treatment, 451

Rural Agricultural Area
Potential assessment of establishing a renewable energy plant in a rural agricultural area, 662

Sampling Units
Full-scale biofilter reduction efficiencies assessed using portable 24-hour sampling units, 170

Sea
Energy supplies and future engines for land, sea, and air, 607, 1233

Season
Quantification of personal exposure concentrations to gasoline vehicle emissions in high-end exposure microenvironments: Effects of fuel and season, 1346

Sedimentary Basin
Compositions and greenhouse gas emission factors of flared and vented gas in the Western Canadian Sedimentary Basin, 992

Selenium Glass Wastes
Feasibility study on cross-linked biopolymeric concrete encapsulating selenium glass wastes, 898

Semiconductor Manufacturing Industries
Integration of a nonmetallic electrostatic precipitator and a wet scrubber for improved removal of particles and corrosive gas cleaning in semiconductor manufacturing industries, 905

Sensitivity and Linearity Analysis
Sensitivity and linearity analysis of ozone in East Asia: The effects of domestic emission and intercontinental transport, 1102

Sewage Sludge
Resource recycling through artificial lightweight aggregates from sewage sludge and derived ash using boric acid flux to lower co-melting temperature, 262

Shale Gas Development
Effect of biogas generation on radon emissions from landfills receiving radium-bearing waste from shale gas development, 1040

Silica
Development of regenerative dye impregnated mesoporous silica materials for assessing exposure to ammonia, 838

Silt Loading
A study on the characteristics of silt loading on paved roads in the Seoul metropolitan area using a mobile monitoring system, 846

Sintering Atmosphere
Effects of sintering atmosphere on cement clinkers produced from chromium-bearing sludge, 587

Sludge
Co-melting technology in resource recycling of sludge derived from stone processing, 1449

SO_2
Effect of solution pH on SO_2, NOx, and Hg removal from simulated coal combustion flue gas in an oxidant-enhanced wet scrubber, 212

Experimental study on removals of SO_2 and NO_x using adsorption of activated carbon/microwave desorption, 1012

NO_2 and SO_2 dispersion modeling and relative roles of emission sources over Map Ta Phut industrial area, Thailand, 932

Sodium Chlorite
The importance of the location of sodium chlorite application in a multipollutant flue gas cleaning system, 707

Soil Amendment
Mercury emission and plant uptake of trace elements during early stage of soil amendment using flue gas desulfurization materials, 139

Source Apportionment
Multiple-year black carbon measurements and source apportionment using Delta-C in Rochester, New York, 880

Source Attribution
Source attribution for mercury deposition in the contiguous United States: Regional difference and seasonal variation, 52

Source Emission Factors
Simulating industrial emissions using Atmospheric Dispersion Modeling System: Model performance and source emission factors, 336

Spatial Distribution of Nitrogen Dioxide
Assessing the spatial distribution of nitrogen dioxide in London, Ontario, 1335

Speciated Arsenic in Air
Speciated arsenic in air: Measurement methodology and risk assessment considerations, 2

Spraying
Rapidly eliminating pathogenic microorganisms in large air space using spraying •OH radicals, 393

State Implementation Plans
Efficacy of recent state implementation plans for 8-hour ozone, 252

Stone Processing
Co-melting technology in resource recycling of sludge derived from stone processing, 1449

Street Sweeping Routes
A novel methodology for determining low-cost fine particulate matter street sweeping routes, 242

Subway Stations
A real-time monitoring and assessment method for calculation of total amounts of indoor air pollutants emitted in subway stations, 517

Sulfur-Modified Activated Carbons
Enhanced mercuric chloride adsorption onto sulfur-modified activated carbons derived from waste tires, 799

Summertime
Evaluating the effects of climate change on summertime ozone using a relative response factor approach for policymakers, 1061

Tapered Element Oscillating Microbalance
Field evaluation of particulate matter measurements using tapered element oscillating microbalance in a layer house, 322

Term Birth Weight
Ambient air pollution and term birth weight in Texas from 1998 to 2004, 1285

Thermal Inactivation
Thermal inactivation of *Bacillus anthracis* surrogate spores in a bench-scale enclosed landfill gas flare, 151

Thermal Processing
Effect of thermal processing on the characteristics of incineration fly ash, 500

***Thiomonas* sp.**
Middle-thermophilic sulfur-oxidizing bacteria *Thiomonas* sp. RAN5 strain for hydrogen sulfide removal, 38

TiO_2
Gas-phase elemental mercury removal in a simulated combustion flue gas using TiO_2 with fluorescent light, 1208

TiO_2 Catalysts
Effect of the Mn oxidation state and lattice oxygen in Mn-based TiO_2 catalysts on the low-temperature selective catalytic reduction of NO by NH_3, 1085

Toluene
Adsorption Behavior of Toluene on Modified 1X Molecular Sieves, 1227

Transformation of microflora during degradation of gaseous toluene in a biofilter detected using PCR-DGGE, 748

Toluene Vapor
Toluene vapor capture by activated carbon particles in a dual gas-solid cyclone system, 924

Toluene Vapor Treatment
Optimizing nutrient supply in a rotatory-switching biofilter for toluene vapor treatment, 451

Trace Elements
Mercury emission and plant uptake of trace elements during early stage of soil amendment using flue gas desulfurization materials, 139

Tracer Gas
Near-road multipollutant profiles: Associations between volatile organic compounds and a tracer gas surrogate near a busy highway, 594

Tracer Gas Surrogate
Near-road multipollutant profiles: Associations between volatile organic compounds and a tracer gas surrogate near a busy highway, 594

Transit Buses
Comparison of on-road emissions for B-0, B-10, and B-20 in transit buses, 443

2,2,4-Trimethyl, 1,3-Pentanediol Monoisobutyrate
Ambient measurements of 2,2,4-trimethyl, 1,3-pentanediol monoisobutyrate in Southern California, 680

Tropospheric Ozone Levels
Extreme value modeling for the analysis and prediction of time series of extreme tropospheric ozone levels: A case study, 651

Tunnel Study
A tunnel study to estimate emission factors from mobile sources in Monterrey, Mexico, 1431

Ultrafine Particles
Exposure to airborne ultrafine particles from cooking in Portuguese homes, 1116

Ultrafine Particle Emission
Ultrafine particle emission from incinerators: The role of the fabric filter, 103

Ultrafine Secondary Organic Aerosols
Enhancement of the deposition of ultrafine secondary organic aerosols by the negative air ion and the effect of relative humidity, 1296

Ultraviolet Light-Emitting Diode Photocatalysis
Application of ultraviolet light-emitting diode photocatalysis to remove volatile organic compounds from indoor air, 1032

Uncertainty
A generalized fuzzy linear programming approach for environmental management problem under uncertainty, 72

Unidirectional-Flow Biotrickling Filters
Comparison of the purification performance and microbial community functional diversity in flowdirectional- switching and unidirectional-flow biotrickling filters, 1203

Urban Environment
Assessment of exposure to airborne ultrafine particles in the urban environment of Lisbon, Portugal, 373

Vehicle Emissions
Multispecies remote sensing measurements of vehicle emissions on Sherman Way in Van Nuys, California, 1127

Vehicle Exhaust Pollutants
Experimental studies of the dilution of vehicle exhaust pollutants by environment-protecting pervious pavement, 92

Vehicular Exhausts Sources
Carbonyl emissions from vehicular exhausts sources in Hong Kong, 221

Volatile Organic Compounds (VOCs)
Application of ultraviolet light-emitting diode photocatalysis to remove volatile organic compounds from indoor air, 1032

Effects of activated carbon surface properties on the adsorption of volatile organic compounds, 1196

The emission patterns of volatile organic compounds during aerobic biotreatment of municipal solid waste using continuous and intermittent aeration, 461

Modeling an air pollution episode in northwestern United States: Identifying the effect of nitrogen oxide and volatile organic compound emission changes on air pollutants formation using direct sensitivity analysis, 1150

Near-road multipollutant profiles: Associations between volatile organic compounds and a tracer gas surrogate near a busy highway, 594

Volatile organic compound emissions from municipal solid waste disposal sites: A case study of Mumbai, India, 398

Waste Combustions
Atmospheric mercury emissions from waste combustions measured by continuous monitoring devices, 686

Waste Tires
Enhanced mercuric chloride adsorption onto sulfur-modified activated carbons derived from waste tires, 799

Wastewater Treatment Plant
Characterization of the olfactory impact around a wastewater treatment plant: Optimization and validation of a hydrogen sulfide determination procedure based on passive diffusion sampling, 863

Estimation of hydrogen sulfide emission rates at several wastewater treatment plants through experimental concentration measurements and dispersion modeling, 758

Wet Scrubber
Integration of a nonmetallic electrostatic precipitator and a wet scrubber for improved removal of particles and corrosive gas cleaning in semiconductor manufacturing industries, 905

Wind Tunnel Test
A wind tunnel test of newly developed personal bioaerosol samplers, 828

Wood Combustion
Obtaining polycyclic aromatic hydrocarbon concentration ratios and molecular markers for residential wood combustion: Temuco, a case study, 44

Xytetracycline Bacterial Residue
Preparation, characterization, and phenol adsorption of activated carbons from oxytetracycline bacterial residue, 1394

Author Index

Abi-Esber, L.
Simulating industrial emissions using Atmospheric Dispersion Modeling System: Model performance and source emission factors, 336

Abichou, T.
A new approach to characterize emission contributions from area sources during optical remote sensing technique testing, 1403
Methane emissions from 20 landfills across the United States using vertical radial plume mapping, 183

Abraham, R.G.
Evaluating the effects of climate change on summertime ozone using a relative response factor approach for policymakers, 1061

Akdeniz, N.
Full-scale biofilter reduction efficiencies assessed using portable 24-hour sampling units, 170

Alberini, A.
Cancer values of prevented fatalities (VPFs), one size does not fit all: The benefits of contaminated site cleanups in Italy, 783

Albuquerque, P.C.
Assessment of exposure to airborne ultrafine particles in the urban environment of Lisbon, Portugal, 373
Exposure to airborne ultrafine particles from cooking in Portuguese homes, 1116

Alonso, M.
Influence of particle location on the number of charges per charged nanoparticle at the outlet of a needle charger, 87

Amar, P.
Development of risk-based air quality management strategies under impacts of climate change, 557

Ana, G.R.E.E.
Polycyclic aromatic hydrocarbon burden in ambient air in selected Niger Delta communities in Nigeria, 18

Anderson, B.E.
Determination of the emissions from an aircraft auxiliary power unit (APU) during the Alternative Aviation Fuel Experiment (AAFEX), 420

Araizaga, A.E.
A tunnel study to estimate emission factors from mobile sources in Monterrey, Mexico, 1431

Arisoy, M.
Multiple comparisons of organic, microbial, and fine particulate pollutants in typical indoor environments: Diurnal and seasonal variation, 1380

Asano, R.
Middle-thermophilic sulfur-oxidizing bacteria *Thiomonas* sp. RAN5 strain for hydrogen sulfide removal, 38

Auvermann, B.W.
Particulate matter emission rates from beef cattle feedlots in Kansas—Reverse dispersion modeling, 350

Avise, J.
Evaluating the effects of climate change on summertime ozone using a relative response factor approach for policymakers, 1061

Bai, H.
Development of regenerative dye impregnated mesoporous silica materials for assessing exposure to ammonia, 838

Bai, M.
Rapidly eliminating pathogenic microorganisms in large air space using spraying •OH radicals, 393

Balachandran, S.
Development of outcome-based, multipollutant mobile source indicators, 431

Baldauf, R.W.
Near-road multipollutant profiles: Associations between volatile organic compounds and a tracer gas surrogate near a busy highway, 594

Balducci, D.
Temporal and spatial variations of polycyclic aromatic hydrocarbon concentrations around a coke oven plant, 1003

Bandemehr, A.
Demonstration of fuel switching on oceangoing vessels in the Gulf of Mexico, 1093

Barker, D.E.
Comparison of six particle size distribution models on the goodness-of-fit to particulate matter sampled from animal buildings, 725

Barlaz, M.
A new approach to characterize emission contributions from area sources during optical remote sensing technique testing, 1403

Barnes, J.E.
An inexpensive active optical remote sensing instrument for assessing aerosol distributions, 198

Barth, E.F.
Evaluation of a sustainable remediation option: Beneficial reuse of petroleum-contaminated sediment as an energy source, 1277

Barzyk, T.M.
Near-road multipollutant profiles: Associations between volatile organic compounds and a tracer gas surrogate near a busy highway, 594

Beghelli, A.
A novel methodology for determining low-cost fine particulate matter street sweeping routes, 242

Belal, E.B.
Effects of foliar application with compost tea and filtrate biogas slurry liquid on yield and fruit quality of Washington navel orange *(Citrus sinenesis Osbeck)* trees, 767

Bell, M.L.
Ambient air pollution and term birth weight in Texas from 1998 to 2004, 1285

Benke, R.R.
Effect of biogas generation on radon emissions from landfills receiving radium-bearing waste from shale gas development, 1040

Beyersdorf, A.J.
Determination of the emissions from an aircraft auxiliary power unit (APU) during the Alternative Aviation Fuel Experiment (AAFEX), 420

Bharadwaj, H.
Evaluation of a sustainable remediation option: Beneficial reuse of petroleum-contaminated sediment as an energy source, 1277

Billo, R.
Air impacts from three alternatives for producing JP-8 jet fuel, 1182
Analysis and comparison of inertinite-derived adsorbent with conventional adsorbents, 489

Bishop, G.A.
Comparison of the MOVES2010a, MOBILE6.2, and EMFAC2007 mobile source emission models with on-road traffic tunnel and remote sensing measurements, 1134
Multispecies remote sensing measurements of vehicle emissions on ShermanWay in Van Nuys, California, 1127

Blazquez, C.A.
A novel methodology for determining low-cost fine particulate matter street sweeping routes, 242

Bonifacio, H.F.
Particulate matter emission rates from beef cattle feedlots in Kansas—Reverse dispersion modeling, 350

Bonnet, S.
Ozone and its potential control strategy for Chon Buri city, Thailand, 1411

Bordado, J.C.
Assessment of exposure to airborne ultrafine particles in the urban environment of Lisbon, Portugal, 373
Exposure to airborne ultrafine particles from cooking in Portuguese homes, 1116

Borhan, M.S.
Comparison of seasonal phenol and p-cresol emissions from ground-level area sources in a dairy operation in central Texas, 381

Bregendahl, K.
Reducing ammonia emissions from laying-hen houses through dietary manipulation, 160

Breysse, P.N.
Assessment of heterogeneity of metal composition of fine particulate matter collected from eight U.S. counties using principal component analysis, 773

Brook, J.R.
Assessing the spatial distribution of nitrogen dioxide in London, Ontario, 1335

Browning, L.
Demonstration of fuel switching on oceangoing vessels in the Gulf of Mexico, 1093

Bufalino, C.
Ambient measurements of 2,2,4-trimethyl, 1,3-pentanediol monoisobutyrate in Southern California, 680

Bujalski, N.M.
Evaluation of a sustainable remediation option: Beneficial reuse of petroleum-contaminated sediment as an energy source, 1277

Bulzan, D.I.
Determination of the emissions from an aircraft auxiliary power unit (APU) during the Alternative Aviation Fuel Experiment (AAFEX), 420

Bumiller, K.
Ambient measurements of 2,2,4-trimethyl, 1,3-pentanediol monoisobutyrate in Southern California, 680

Buonanno, G.
Ultrafine particle emission from incinerators: The role of the fabric filter, 103

Burns, R.T.
Reducing ammonia emissions from laying-hen houses through dietary manipulation, 160

Büyüksönmez, F.
Mitigation of odor causing emissions—Bench-scale investigation, 1423

Buzzelli, M.
Assessing the spatial distribution of nitrogen dioxide in London, Ontario, 1335

Caballero, P.
Chemical characterization and factor analysis of $PM_{2.5}$ in two sites of Monterrey, Mexico, 817

Campbell, D.
Quantification of personal exposure concentrations to gasoline vehicle emissions in high-end exposure microenvironments: Effects of fuel and season, 1346

Campbell, D.E.
Comparison of the MOVES2010a, MOBILE6.2, and EMFAC2007 mobile source emission models with on-road traffic tunnel and remote sensing measurements, 1134

Campleman, S.L.
Speciated arsenic in air: Measurement methodology and risk assessment considerations, 2

Campos Candel, A.
Characterization of the olfactory impact around a wastewater treatment plant: Optimization and validation of a hydrogen sulfide determination procedure based on passive diffusion sampling, 863

Cao, J.-J.
Winter and Summer $PM_{2.5}$ Chemical Compositions in Fourteen Chinese Cities, 1214

Cao, W.
Application of neutral electrolyzed water spray for reducing dust levels in a layer breeding house, 1329

Capareda, S.
Comparison of seasonal phenol and p-cresol emissions from ground-level area sources in a dairy operation in central Texas, 381

Carrillo, O.
Chemical characterization and factor analysis of $PM_{2.5}$ in two sites of Monterrey, Mexico, 817

Cen, K.-F.
Dioxins and their fingerprint in size-classified fly ash fractions from municipal solid waste incinerators in China—Mechanical grate and fluidized bed units, 717

Ceppi, M.
Temporal and spatial variations of polycyclic aromatic hydrocarbon concentrations around a coke oven plant, 1003

Cereceda-Balic, F.
Obtaining polycyclic aromatic hydrocarbon concentration ratios and molecular markers for residential wood combustion: Temuco, a case study, 44

Chai, F.
Control strategies of atmospheric mercury emissions from coal-fired power plants in China, 576

Chalupa, D.C.
Multiple-year black carbon measurements and source apportionment using Delta-C in Rochester, New York, 880

Chang, J.-E.
Effects of sintering atmosphere on cement clinkers produced from chromium-bearing sludge, 587

Chang, S.-Y.
Evaluating the performance of the horizontal radial plume mapping technique for locating multiple plumes, 1249

Chang, Y.-C.
Development of regenerative dye impregnated mesoporous silica materials for assessing exposure to ammonia, 838

Chang, Y.-J.
An economic evaluation and assessment of environmental impact of the municipal solid waste management system for Taichung City in Taiwan, 527

Chang, Y.-N.
Mercury emission and plant uptake of trace elements during early stage of soil amendment using flue gas desulfurization materials, 139

Chanton, J.
A new approach to characterize emission contributions from area sources during optical remote sensing technique testing, 1403

Methane emissions from 20 landfills across the United States using vertical radial plume mapping, 183

Chen, J.
Evaluating the effects of climate change on summertime ozone using a relative response factor approach for policymakers, 1061

Chen, J.-W.
Experimental studies of the dilution of vehicle exhaust pollutants by environment-protecting pervious pavement, 92

Chen, L.W.A.
Fine particulate matter and visibility in the Lake Tahoe Basin: Chemical characterization, trends, and source apportionment, 953

Chen, Q.
Effectiveness and mechanisms of hydrogen sulfide adsorption by camphor-derived biochar, 873

Chen, T.
Hydrothermal treatment for inactivating some hygienic microbial indicators from food waste–amended animal feed, 810

Chen, T.-H.
Experimental studies of the dilution of vehicle exhaust pollutants by environment-protecting pervious pavement, 92

Chen, W.-J.
Enhanced mercuric chloride adsorption onto sulfur-modified activated carbons derived from waste tires, 799

Chen, Y.-L.
Effects of sintering atmosphere on cement clinkers produced from chromium-bearing sludge, 587

Cheng, C.-M.
Mercury emission and plant uptake of trace elements during early stage of soil amendment using flue gas desulfurization materials, 139

Cheng, K.
Control strategies of atmospheric mercury emissions from coal-fired power plants in China, 576

Cheng, W.-H.
Breakthrough indicator for aromatic VOCs using needle trap samplers for activated carbon adsorbent, 946

Cheng, Y.
Carbonyl emissions from vehicular exhausts sources in Hong Kong, 221

Cheng, Y.S.
Awind tunnel test of newly developed personal bioaerosol samplers, 828

Cheung, K.
Historical trends in the mass and chemical species concentrations of coarse particulate matter in the Los Angeles Basin and relation to sources and air quality regulations, 541

Chi, Z.
Effect of biocover equipped with a novel passive air diffusion system on microbial methane oxidation and community of methanotrophs, 278

Chiang, H.-S.
Development of regenerative dye impregnated mesoporous silica materials for assessing exposure to ammonia, 838

Cho, J.H.
Gas-phase elemental mercury removal in a simulated combustion flue gas using TiO_2 with fluorescent light, 1208

Chow, J.C.
Comparison of the MOVES2010a, MOBILE6.2, and EMFAC2007 mobile source emission models with on-road traffic tunnel and remote sensing measurements, 1134

Energy supplies and future engines for land, sea, and air, 1233

Winter and Summer $PM_{2.5}$ Chemical Compositions in Fourteen Chinese Cities, 1214

Chu, C.-W.
An economic evaluation and assessment of environmental impact of the municipal solid waste management system for Taichung City in Taiwan, 527

Chu, H.-W.
Source attribution for mercury deposition in the contiguous United States: Regional difference and seasonal variation, 52

Chudnovsky, A.A.
Prediction of daily fine particulate matter concentrations using aerosol optical depth retrievals from the Geostationary Operational Environmental Satellite (GOES), 1022

Chuen, O.C.
Benefits of clean development mechanism application on the life cycle assessment perspective: A case study in the palm oil industry, 299

Chung, S.H.
Evaluating the effects of climate change on summertime ozone using a relative response factor approach for policymakers, 1061

Chusai, C.
NO_2 and SO_2 dispersion modeling and relative roles of emission sources over Map Ta Phut industrial area, Thailand, 932

Ciesielski, A.
Near-road multipollutant profiles: Associations between volatile organic compounds and a tracer gas surrogate near a busy highway, 594

Cipolla, M.
Temporal and spatial variations of polycyclic aromatic hydrocarbon concentrations around a coke oven plant, 1003

Clark, C.
The evolution and impact of testing baghouse filter performance, 916

Clark, J.
A new approach to characterize emission contributions from area sources during optical remote sensing technique testing, 1403

Clark, P.J.
Evaluation of a sustainable remediation option: Beneficial reuse of petroleum-contaminated sediment as an energy source, 1277

Clifford, E.
Optimization of a horizontal-flow biofilm reactor for the removal of methane at low temperatures, 1166

Clifford, J.M.
A 2009 Mobile Source Carbon Dioxide Emissions Inventory for the University of Central Florida, 1050

Cocker III, D.R.
Aging of secondary organic aerosol from α-pinene ozonolysis: Roles of hydroxyl and nitrate radicals, 1359

Cocker, K.
Ambient measurements of 2,2,4-trimethyl, 1,3-pentanediol monoisobutyrate in Southern California, 680

Coderre, A.R.
Compositions and greenhouse gas emission factors of flared and vented gas in the Western Canadian Sedimentary Basin, 992

Cohan, D.S.
Efficacy of recent state implementation plans for 8-hour ozone, 252

Collins, G.
Optimization of a horizontal-flow biofilm reactor for the removal of methane at low temperatures, 1166

Coll Lozano, C.
Characterization of the olfactory impact around a wastewater treatment plant: Optimization and validation of a hydrogen sulfide determination procedure based on passive diffusion sampling, 863

Cooper, C.D.
A 2009 Mobile Source Carbon Dioxide Emissions Inventory for the University of Central Florida, 1050

Corsi, R.L.
Emissions of *p*-dichlorobenzene and naphthalene from consumer products, 1075

Costagliola, M.A.
Bioethanol–gasoline fuel blends: Exhaust emissions and morphological characterization of particulate from a moped engine, 888

Cotton, M.
Mitigation of odor causing emissions—Bench-scale investigation, 1423

Couzo, E.
Assessment of a regulatory model's performance relative to large spatial heterogeneity in observed ozone in Houston, Texas, 696

Darrow, L.A.
Development of outcome-based, multipollutant mobile source indicators, 431

DenBleyker, A.
Comparison of the MOVES2010a, MOBILE6.2, and EMFAC2007 mobile source emission models with on-road traffic tunnel and remote sensing measurements, 1134

Effects of improved spatial and temporal modeling of on-road vehicle emissions, 471

Dennis, B.H.
Air impacts from three alternatives for producing JP-8 jet fuel, 1182

Analysis and comparison of inertinite-derived adsorbent with conventional adsorbents, 489

Díaz-Robles, L.A.
Obtaining polycyclic aromatic hydrocarbon concentration ratios and molecular markers for residential wood combustion: Temuco, a case study, 44

Dietze, V.
A comparative study on the varying exposure to atmospheric fine and coarse particles under urban and rural conditions, 1313

Digar, A.
Efficacy of recent state implementation plans for 8-hour ozone, 252

Domínguez, A.M.
Obtaining polycyclic aromatic hydrocarbon concentration ratios and molecular markers for residential wood combustion: Temuco, a case study, 44

Dong, J.-H.
Mercury emission and plant uptake of trace elements during early stage of soil amendment using flue gas desulfurization materials, 139

Dong, X.
Sensitivity and linearity analysis of ozone in East Asia: The effects of domestic emission and intercontinental transport, 1102

Douglass, C.
Efficacy of recent state implementation plans for 8-hour ozone, 252

Du, Y.
Dioxins and their fingerprint in size-classified fly ash fractions from municipal solid waste incinerators in China—Mechanical grate and fluidized bed units, 717

Duan, E.
Characteristics of penicillin bacterial residue, 485
Preparation, characterization, and phenol adsorption of activated carbons from oxytetracycline bacterial residue, 1394

Duan, Z.
Effect of biocover equipped with a novel passive air diffusion system on microbial methane oxidation and community of methanotrophs, 278

DuBois, D.W.
Fine particulate matter and visibility in the Lake Tahoe Basin: Chemical characterization, trends, and source apportionment, 953

Duhl, T.
Evaluating the effects of climate change on summertime ozone using a relative response factor approach for policymakers, 1061

El-Abd, A.El-N.A.
Effects of foliar application with compost tea and filtrate biogas slurry liquid on yield and fruit quality of Washington navel orange *(Citrus sinenesis Osbeck)* trees, 767

El-Fadel, M.
Simulating industrial emissions using Atmospheric Dispersion Modeling System: Model performance and source emission factors, 336

Emerole, G.O.
Polycyclic aromatic hydrocarbon burden in ambient air in selected Niger Delta communities in Nigeria, 18

Endlicher, W.
A comparative study on the varying exposure to atmospheric fine and coarse particles under urban and rural conditions, 1313

England, G.C.
Energy supplies and future engines for land, sea, and air, 1233

Eom, Y.
Gas-phase elemental mercury removal in a simulated combustion flue gas using TiO_2 with fluorescent light, 1208

Escarela, G.
Extreme value modeling for the analysis and prediction of time series of extreme tropospheric ozone levels: A case study, 651

Espinós Morató, H.
Characterization of the olfactory impact around a wastewater treatment plant: Optimization and validation of a hydrogen sulfide determination procedure based on passive diffusion sampling, 863
Estimation of hydrogen sulfide emission rates at several wastewater treatment plants through experimental concentration measurements and dispersion modeling, 758

Etcharren, P.
Obtaining polycyclic aromatic hydrocarbon concentration ratios and molecular markers for residential wood combustion: Temuco, a case study, 44

Fadic, X.
Obtaining polycyclic aromatic hydrocarbon concentration ratios and molecular markers for residential wood combustion: Temuco, a case study, 44

Fan, Y.
A generalized fuzzy linear programming approach for environmental management problem under uncertainty, 72

Fang, J.-J.
The emission patterns of volatile organic compounds during aerobic biotreatment of municipal solid waste using continuous and intermittent aeration, 461

Faulkner, W.B.
Comparison of seasonal phenol and p-cresol emissions from ground-level area sources in a dairy operation in central Texas, 381

Feng, N.S.Y.
Carbonyl emissions from vehicular exhausts sources in Hong Kong, 221

Fitz, D.R.
Ambient measurements of 2,2,4-trimethyl, 1,3-pentanediol monoisobutyrate in Southern California, 680

Fu, J.S.
Sensitivity and linearity analysis of ozone in East Asia: The effects of domestic emission and intercontinental transport, 1102

Fu, Y.-P.
Co-melting technology in resource recycling of sludge derived from stone processing, 1449
Resource recycling through artificial lightweight aggregates from sewage sludge and derived ash using boric acid flux to lower co-melting temperature, 262

Fujita, E.
Quantification of personal exposure concentrations to gasoline vehicle emissions in high-end exposure microenvironments: Effects of fuel and season, 1346

Fujita, E.M.
Comparison of the MOVES2010a, MOBILE6.2, and EMFAC2007 mobile source emission models with on-road traffic tunnel and remote sensing measurements, 1134

Gangupomu, R.H.
Air impacts from three alternatives for producing JP-8 jet fuel, 1182
Analysis and comparison of inertinite-derived adsorbent with conventional adsorbents, 489

Gao, L.
Experimental study on removals of SO_2 and NO_x using adsorption of activated carbon/microwave desorption, 1012

Gao, Q.
Preparation, characterization, and phenol adsorption of activated carbons from oxytetracycline bacterial residue, 1394

Gao, Y.
Sensitivity and linearity analysis of ozone in East Asia: The effects of domestic emission and intercontinental transport, 1102

Gao, Z.
Determination of partition and diffusion coefficients of formaldehyde in selected building materials and impact of relative humidity, 671

Gathright, K.
Demonstration of fuel switching on oceangoing vessels in the Gulf of Mexico, 1093

Geer, L.A.
Ambient air pollution and term birth weight in Texas from 1998 to 2004, 1285

Gerrity, S.
Optimization of a horizontal-flow biofilm reactor for the removal of methane at low temperatures, 1166

Geyh, A.S.
Assessment of heterogeneity of metal composition of fine particulate matter collected from eight U.S. counties using principal component analysis, 773

Goldsmith, D.
A new approach to characterize emission contributions from area sources during optical remote sensing technique testing, 1403

Goldsmith, Jr., C.D.
Methane emissions from 20 landfills across the United States using vertical radial plume mapping, 183

Goliff, W.S.
Ambient measurements of 2,2,4-trimethyl, 1,3-pentanediol monoisobutyrate in Southern California, 680

Gomes, J.F.
Assessment of exposure to airborne ultrafine particles in the urban environment of Lisbon, Portugal, 373
Exposure to airborne ultrafine particles from cooking in Portuguese homes, 1116

Gómez Pérez, T.
Characterization of the olfactory impact around a wastewater treatment plant: Optimization and validation of a hydrogen sulfide determination procedure based on passive diffusion sampling, 863

Gong, L.
Characteristics of penicillin bacterial residue, 485

Green, M.C.
Fine particulate matter and visibility in the Lake Tahoe Basin: Chemical characterization, trends, and source apportionment, 953

Green, R.
A new approach to characterize emission contributions from area sources during optical remote sensing technique testing, 1403
Methane emissions from 20 landfills across the United States using vertical radial plume mapping, 183

Gu, W.
The addition of modified attapulgite reduces the emission of nitrous oxide and ammonia from aerobically composted chicken manure, 1174

Guenther, A.
Evaluating the effects of climate change on summertime ozone using a relative response factor approach for policymakers, 1061

Guerrero, P.A.
Emissions of *p*-dichlorobenzene and naphthalene from consumer products, 1075

Guevara, J.L.
Obtaining polycyclic aromatic hydrocarbon concentration ratios and molecular markers for residential wood combustion: Temuco, a case study, 44

Guignet, D.
Cancer values of prevented fatalities (VPFs), one size does not fit all: The benefits of contaminated site cleanups in Italy, 783

Güllü, G.
Multiple comparisons of organic, microbial, and fine particulate pollutants in typical indoor environments: Diurnal and seasonal variation, 1380

Guo, B.
Characteristics of penicillin bacterial residue, 485
Preparation, characterization, and phenol adsorption of activated carbons from oxytetracycline bacterial residue, 1394

Ha, J.-H.
Emissions monitoring at a deep-pit swine finishing facility: Research methods and system performance, 1264

Hagen, D.
Determination of the emissions from an aircraft auxiliary power unit (APU) during the Alternative Aviation Fuel Experiment (AAFEX), 420

Hallmark, S.
Comparison of on-road emissions for B-0, B-10, and B-20 in transit buses, 443

Han, B.
Integration of a nonmetallic electrostatic precipitator and a wet scrubber for improved removal of particles and corrosive gas cleaning in semiconductor manufacturing industries, 905

Han, I.
Assessment of heterogeneity of metal composition of fine particulate matter collected from eight U.S. counties using principal component analysis, 773

Han, J.
Characteristics of penicillin bacterial residue, 485

Han, S.
A study on the characteristics of silt loading on paved roads in the Seoul metropolitan area using a mobile monitoring system, 846

Han, Y.-M.
Winter and Summer $PM_{2.5}$ Chemical Compositions in Fourteen Chinese Cities, 1214

Hao, J.
Control strategies of atmospheric mercury emissions from coal-fired power plants in China, 576

Happonen, M.
Study of Miller timing on exhaust emissions of a hydrotreated vegetable oil (HVO)-fueled diesel engine, 1305

Hara, K.
Driver exposure to particulate matter in Bangkok, 64

Harner, III, J.P.
Particulate matter emission rates from beef cattle feedlots in Kansas—Reverse dispersion modeling, 350

Hartley, S.
Demonstration of fuel switching on oceangoing vessels in the Gulf of Mexico, 1093

Hasegawa, S.
Driver exposure to particulate matter in Bangkok, 64

Hater, G.
A new approach to characterize emission contributions from area sources during optical remote sensing technique testing, 1403
Methane emissions from 20 landfills across the United States using vertical radial plume mapping, 183

He, P.-J.
The emission patterns of volatile organic compounds during aerobic biotreatment of municipal solid waste using continuous and intermittent aeration, 461

He, S.
Comparison of the purification performance and microbial community functional diversity in flowdirectional-switching and unidirectional-flow biotrickling filters, 1203

He, X.
The emission patterns of volatile organic compounds during aerobic biotreatment of municipal solid waste using continuous and intermittent aeration, 461

Heber, A.J.
Emissions monitoring at a deep-pit swine finishing facility: Research methods and system performance, 1264
Field evaluation of particulate matter measurements using tapered element oscillating microbalance in a layer house, 322

Heikkilä, J.
Study of Miller timing on exhaust emissions of a hydrotreated vegetable oil (HVO)-fueled diesel engine, 1305

Herndon, S.C.
Determination of the emissions from an aircraft auxiliary power unit (APU) during the Alternative Aviation Fuel Experiment (AAFEX), 420

Hidy, G.M.
Energy supplies and future engines for land, sea, and air, 1233

Higuchi, T.
Optimizing nutrient supply in a rotatory-switching biofilter for toluene vapor treatment, 451
Transformation ofmicroflora during degradation of gaseous toluene in a biofilter detected using PCR-DGGE, 748

Hirooka, K.
Middle-thermophilic sulfur-oxidizing bacteria *Thiomonas* sp. RAN5 strain for hydrogen sulfide removal, 38

Hixson, M.
Resolving the interactions between population density and air pollution emissions controls in the San Joaquin Valley, USA, 566

Ho, K.F.
Carbonyl emissions from vehicular exhausts sources in Hong Kong, 221
Winter and Summer $PM_{2.5}$ Chemical Compositions in Fourteen Chinese Cities, 1214

Ho, S.S.H.
Carbonyl emissions from vehicular exhausts sources in Hong Kong, 221

Hong, S.C.
Effect of the Mn oxidation state and lattice oxygen in Mn-based TiO_2 catalysts on the low-temperature selective catalytic reduction of NO by NH_3, 1085
Improving the activity of Mn/TiO2 catalysts through control of the pH and valence state of Mn during their preparation, 362
Physicochemical properties of chars at different treatment temperatures, 235

Hopke, P.K.
Multiple-year black carbon measurements and source apportionment using Delta-C in Rochester, New York, 880

Howard, R.
Determination of the emissions from an aircraft auxiliary power unit (APU) during the Alternative Aviation Fuel Experiment (AAFEX), 420

Hsu, J.-N.
Development of regenerative dye impregnated mesoporous silica materials for assessing exposure to ammonia, 838

Hu, C.-C.
Mercury emission and plant uptake of trace elements during early stage of soil amendment using flue gas desulfurization materials, 139

Hu, J.
Resolving the interactions between population density and air pollution emissions controls in the San Joaquin Valley, USA, 566

Hu, S.-C.
Co-melting technology in resource recycling of sludge derived from stone processing, 1449
Resource recycling through artificial lightweight aggregates from sewage sludge and derived ash using boric acid flux to lower co-melting temperature, 262

Hu, S.-H.
Co-melting technology in resource recycling of sludge derived from stone processing, 1449
Resource recycling through artificial lightweight aggregates from sewage sludge and derived ash using boric acid flux to lower co-melting temperature, 262

Hu, Y.
Development of outcome-based, multipollutant mobile source indicators, 431
Modeling an air pollution episode in northwestern United States: Identifying the effect of nitrogen oxide and volatile organic compound emission changes on air pollutants formation using direct sensitivity analysis, 1150

Huang, C.-H.
Influence of particle location on the number of charges per charged nanoparticle at the outlet of a needle charger, 87

Huang, G.
A generalized fuzzy linear programming approach for environmental management problem under uncertainty, 72

Huang, S.
Removal of nitric oxide in a biotrickling filter under thermophilic condition using *Chelatococcus daeguensis*, 509

Huang, S.-C.
Breakthrough indicator for aromatic VOCs using needle trap samplers for activated carbon adsorbent, 946

Huang, W.-J.
Potential assessment of establishing a renewable energy plant in a rural agricultural area, 662

Huang, X.
The addition of modified attapulgite reduces the emission of nitrous oxide and ammonia from aerobically composted chicken manure, 1174

Huang, Y.
Carbonyl emissions from vehicular exhausts sources in Hong Kong, 221
The effects of manganese precursors on Mn-based/TiO_2 catalysts for catalytic reduction of NO with NH_3, 271

Huang, Y.-N.
Breakthrough indicator for aromatic VOCs using needle trap samplers for activated carbon adsorbent, 946

Hudgins, C.H.
Determination of the emissions from an aircraft auxiliary power unit (APU) during the Alternative Aviation Fuel Experiment (AAFEX), 420

Hunt, Jr., W.
Quantification of emission factor uncertainty, 287

Hutson, N.D.
Effect of solution pH on SO_2, NOx, and Hg removal from simulated coal combustion flue gas in an oxidant-enhanced wet scrubber, 212
The importance of the location of sodium chlorite application in a multipollutant flue gas cleaning system, 707

Ie, I.-R.
Enhanced mercuric chloride adsorption onto sulfur-modified activated carbons derived from waste tires, 799

Inoue, M.
Driver exposure to particulate matter in Bangkok, 64

Isakov, V.
Near-road multipollutant profiles: Associations between volatile organic compounds and a tracer gas surrogate near a busy highway, 594

Jang, C.
Source attribution for mercury deposition in the contiguous United States: Regional difference and seasonal variation, 52

Janni, K.A.
Full-scale biofilter reduction efficiencies assessed using portable 24-hour sampling units, 170

Jeffries, H.E.
Assessment of a regulatory model's performance relative to large spatial heterogeneity in observed ozone in Houston, Texas, 696

Jen, Y.-H.
Enhanced mercuric chloride adsorption onto sulfur-modified activated carbons derived from waste tires, 799

Jia, X.
The addition of modified attapulgite reduces the emission of nitrous oxide and ammonia from aerobically composted chicken manure, 1174

Jiang, J.-R.
Breakthrough indicator for aromatic VOCs using needle trap samplers for activated carbon adsorbent, 946

Jin, Y.
Emissions monitoring at a deep-pit swine finishing facility: Research methods and system performance, 1264
Hydrothermal treatment for inactivating some hygienic microbial indicators from food waste–amended animal feed, 810

Jinsart, W.
Driver exposure to particulate matter in Bangkok, 64

Jo, Y.M.
Influence of pleat geometry on filter cleaning in PTFE/glass composite filter, 1257
Toluene vapor capture by activated carbon particles in a dual gas-solid cyclone system, 924

Johnson, M.
Assessing the spatial distribution of nitrogen dioxide in London, Ontario, 1335

Johnson, M.R.
Black carbon particulate matter emission factors for buoyancy-driven associated gas flares, 307
Compositions and greenhouse gas emission factors of flared and vented gas in the Western Canadian Sedimentary Basin, 992

Jou, C.-J.G.
Microwave-induced nanoscale zero-valent iron degradation of perchloroethylene and pentachlorophenol, 1443

Jung, Y.-W.
A study on the characteristics of silt loading on paved roads in the Seoul metropolitan area using a mobile monitoring system, 846

Kaewmanee, C.
Driver exposure to particulate matter in Bangkok, 64

Kaminski, U.
A comparative study on the varying exposure to atmospheric fine and coarse particles under urban and rural conditions, 1313

Kang, O.Y.
A real-time monitoring and assessment method for calculation of total amounts of indoor air pollutants emitted in subway stations, 517

Kang, Y.S.
Physicochemical properties of chars at different treatment temperatures, 235

Kao, N.-H.
Potential assessment of establishing a renewable energy plant in a rural agricultural area, 662

Karaca, F.
Determination of air quality zones in Turkey, 408

Karita, K.
Driver exposure to particulate matter in Bangkok, 64

Karthik, M.
Development of regenerative dye impregnated mesoporous silica materials for assessing exposure to ammonia, 838

Keener, T.C.
Effects of activated carbon surface properties on the adsorption of volatile organic compounds, 1196
Evaluation of a sustainable remediation option: Beneficial reuse of petroleum-contaminated sediment as an energy source, 1277

Kennelly, C.
Optimization of a horizontal-flow biofilm reactor for the removal of methane at low temperatures, 1166

Kerbachi, R.
Measurement of BTEX (benzene, toluene, ethybenzene, and xylene) levels at urban and semirural areas of Algiers City using passive air samplers, 1370

Kerchich, Y.
Measurement of BTEX (benzene, toluene, ethybenzene, and xylene) levels at urban and semirural areas of Algiers City using passive air samplers, 1370

Keskinen, J.
Study of Miller timing on exhaust emissions of a hydrotreated vegetable oil (HVO)-fueled diesel engine, 1305

Kida, A.
Atmospheric mercury emissions from waste combustions measured by continuous monitoring devices, 686

Kim, D.
Feasibility study on cross-linked biopolymeric concrete encapsulating selenium glass wastes, 898

Kim, H.-J.
Integration of a nonmetallic electrostatic precipitator and a wet scrubber for improved removal of particles and corrosive gas cleaning in semiconductor manufacturing industries, 905

Kim, J. K.
Physicochemical properties of chars at different treatment temperatures, 235

Kim, J.T.
A real-time monitoring and assessment method for calculation of total amounts of indoor air pollutants emitted in subway stations, 517

Kim, M.J.
A real-time monitoring and assessment method for calculation of total amounts of indoor air pollutants emitted in subway stations, 517

Kim, S.B.
Influence of pleat geometry on filter cleaning in PTFE/glass composite filter, 1257

Kim, S.S.
Effect of the Mn oxidation state and lattice oxygen in Mn-based TiO_2 catalysts on the low-temperature selective catalytic reduction of NO by NH_3, 1085
Improving the activity of Mn/TiO2 catalysts through control of the pH and valence state of Mn during their preparation, 362
Physicochemical properties of chars at different treatment temperatures, 235

Kim, Y.-J.
Integration of a nonmetallic electrostatic precipitator and a wet scrubber for improved removal of particles and corrosive gas cleaning in semiconductor manufacturing industries, 905

Kinsey, J.S.
Determination of the emissions from an aircraft auxiliary power unit (APU) during the Alternative Aviation Fuel Experiment (AAFEX), 420

Kleeman, M.J.
Resolving the interactions between population density and air pollution emissions controls in the San Joaquin Valley, USA, 566

Kliebenstein, J.
Reducing ammonia emissions from laying-hen houses through dietary manipulation, 160

Knighton, W. B.
Determination of the emissions from an aircraft auxiliary power unit (APU) during the Alternative Aviation Fuel Experiment (AAFEX), 420

Ko, M.-S.
Effects of sintering atmosphere on cement clinkers produced from chromium-bearing sludge, 587

Kositkanawuth, K.
Air impacts from three alternatives for producing JP-8 jet fuel, 1182
Analysis and comparison of inertinite-derived adsorbent with conventional adsorbents, 489

Kostinski, A.
Prediction of daily fine particulate matter concentrations using aerosol optical depth retrievals from the Geostationary Operational Environmental Satellite (GOES), 1022

Kotlov, T.
Prediction of daily fine particulate matter concentrations using aerosol optical depth retrievals from the Geostationary Operational Environmental Satellite (GOES), 1022

Koutrakis, P.
Prediction of daily fine particulate matter concentrations using aerosol optical depth retrievals from the Geostationary Operational Environmental Satellite (GOES), 1022

Krstic, G.
A reanalysis of fine particulate matter air pollution vs. life expectancy in the United States, 989

Krzyzynska, R.
Effect of solution pH on SO_2, NOx, and Hg removal from simulated coal combustion flue gas in an oxidant-enhanced wet scrubber, 212
The importance of the location of sodium chlorite application in a multipollutant flue gas cleaning system, 707

Kwon, D.W.
Effect of the Mn oxidation state and lattice oxygen in Mn-based TiO_2 catalysts on the low-temperature selective catalytic reduction of NO by NH_3, 1085

Lai, Y.-C.
Effects of sintering atmosphere on cement clinkers produced from chromium-bearing sludge, 587

Lam, K.M.
Carbonyl emissions from vehicular exhausts sources in Hong Kong, 221

Lam, Y.F.
Sensitivity and linearity analysis of ozone in East Asia: The effects of domestic emission and intercontinental transport, 1102

Lamb, B.
Evaluating the effects of climate change on summertime ozone using a relative response factor approach for policymakers, 1061

Larmi, M.
Study of Miller timing on exhaust emissions of a hydrotreated vegetable oil (HVO)-fueled diesel engine, 1305

Lawson, D.R.
Comparison of the MOVES2010a, MOBILE6.2, and EMFAC2007 mobile source emission models with on-road traffic tunnel and remote sensing measurements, 1134
Multispecies remote sensing measurements of vehicle emissions on ShermanWay in Van Nuys, California, 1127

Lee, C.-L.
Microwave-induced nanoscale zero-valent iron degradation of perchloroethylene and pentachlorophenol, 1443

Lee, H.D.
Physicochemical properties of chars at different treatment temperatures, 235

Lee, H.J.
Prediction of daily fine particulate matter concentrations using aerosol optical depth retrievals from the Geostationary Operational Environmental Satellite (GOES), 1022

Lee, J.
Comparison of six particle size distribution models on the goodness-of-fit to particulate matter sampled from animal buildings, 725

Lee, J.-Y.
Evaluation of a sustainable remediation option: Beneficial reuse of petroleum-contaminated sediment as an energy source, 1277

Lee, M.-H.
Influence of pleat geometry on filter cleaning in PTFE/glass composite filter, 1257

Lee, S.C.
Winter and Summer $PM_{2.5}$ Chemical Compositions in Fourteen Chinese Cities, 1214
Carbonyl emissions from vehicular exhausts sources in Hong Kong, 221

Lee, S.M.
Effect of the Mn oxidation state and lattice oxygen in Mn-based TiO_2 catalysts on the low-temperature selective catalytic reduction of NO by NH_3, 1085

Lee, T.G.
Gas-phase elemental mercury removal in a simulated combustion flue gas using TiO_2 with fluorescent light, 1208

Legge, A.H.
Energy supplies and future engines for land, sea, and air, 1233

Lehto, K.
Study of Miller timing on exhaust emissions of a hydrotreated vegetable oil (HVO)-fueled diesel engine, 1305

Lewis, A.S.
Speciated arsenic in air: Measurement methodology and risk assessment considerations, 2

Li, B.
Application of neutral electrolyzed water spray for reducing dust levels in a layer breeding house, 1329

Li, H.
Effects of activated carbon surface properties on the adsorption of volatile organic compounds, 1196

Hydrothermal treatment for inactivating some hygienic microbial indicators from food waste–amended animal feed, 810

Reducing ammonia emissions from laying-hen houses through dietary manipulation, 160

Li, J.
An examination of boundary layer structure under the influence of the gap winds in Urumqi, China, during air pollution episode in winter, 26

Li, L.
Effects of activated carbon surface properties on the adsorption of volatile organic compounds, 1196

Li, Q.-F.
Field evaluation of particulate matter measurements using tapered element oscillating microbalance in a layer house, 322

Li, S.
Reducing ammonia emissions from laying-hen houses through dietary manipulation, 160

Li, S.-N.
Development of regenerative dye impregnated mesoporous silica materials for assessing exposure to ammonia, 838

Li, X.
An examination of boundary layer structure under the influence of the gap winds in Urumqi, China, during air pollution episode in winter, 26

Li, X.-D.
Dioxins and their fingerprint in size-classified fly ash fractions from municipal solid waste incinerators in China—Mechanical grate and fluidized bed units, 717

Liang, W.
Removal of nitric oxide in a biotrickling filter under thermophilic condition using *Chelatococcus daeguensis*, 509

Liao, K.-J.
Development of risk-based air quality management strategies under impacts of climate change, 557

Lim, J.J.
A real-time monitoring and assessment method for calculation of total amounts of indoor air pollutants emitted in subway stations, 517

Lim, T.-T.
Emissions monitoring at a deep-pit swine finishing facility: Research methods and system performance, 1264

Lim, Y.H.
Toluene vapor capture by activated carbon particles in a dual gas-solid cyclone system, 924

Limpaseni, W.
Ozone and its potential control strategy for Chon Buri city, Thailand, 1411

Lin, C.
Microwave-induced nanoscale zero-valent iron degradation of perchloroethylene and pentachlorophenol, 1443

Lin, C.-J.
Source attribution for mercury deposition in the contiguous United States: Regional difference and seasonal variation, 52

Lin, C.-W.
Mercury emission and plant uptake of trace elements during early stage of soil amendment using flue gas desulfurization materials, 139

Lin, M.-D.
An economic evaluation and assessment of environmental impact of the municipal solid waste management system for Taichung City in Taiwan, 527

Lin, W.-S.
Experimental studies of the dilution of vehicle exhaust pollutants by environment-protecting pervious pavement, 92

Lindhjem, C.E.
Comparison of the MOVES2010a, MOBILE6.2, and EMFAC2007 mobile source emission models with on-road traffic tunnel and remote sensing measurements, 1134

Effects of improved spatial and temporal modeling of on-road vehicle emissions, 471

Liu, C.-M.
Experimental studies of the dilution of vehicle exhaust pollutants by environment-protecting pervious pavement, 92

Liu, J.
Removal of nitric oxide in a biotrickling filter under thermophilic condition using *Chelatococcus daeguensis*, 509

Liu, R.
Characteristics of penicillin bacterial residue, 485

Liu, X.
Determination of partition and diffusion coefficients of formaldehyde in selected building materials and impact of relative humidity, 671

Liu, Z.
Field evaluation of particulate matter measurements using tapered element oscillating microbalance in a layer house, 322

Llanos, A.L.
Obtaining polycyclic aromatic hydrocarbon concentration ratios and molecular markers for residential wood combustion: Temuco, a case study, 44

Llavador Colomer, F.
Characterization of the olfactory impact around a wastewater treatment plant: Optimization and validation of a hydrogen sulfide determination procedure based on passive diffusion sampling, 863

Estimation of hydrogen sulfide emission rates at several wastewater treatment plants through experimental concentration measurements and dispersion modeling, 758

Lloyd, A.C.
Energy supplies and future engines for land, sea, and air, 1233

Lobo, P.
Determination of the emissions from an aircraft auxiliary power unit (APU) during the Alternative Aviation Fuel Experiment (AAFEX), 420

Long, Y.
Effect of biocover equipped with a novel passive air diffusion system on microbial methane oxidation and community of methanotrophs, 278

Loughlin, D.H.
Evaluating the effects of climate change on summertime ozone using a relative response factor approach for policymakers, 1061

Lu, S.-Y.
Dioxins and their fingerprint in size-classified fly ash fractions from municipal solid waste incinerators in China—Mechanical grate and fluidized bed units, 717

Lu, W.
Effect of biocover equipped with a novel passive air diffusion system on microbial methane oxidation and community of methanotrophs, 278

Lu, W.-C.
Mercury emission and plant uptake of trace elements during early stage of soil amendment using flue gas desulfurization materials, 139

Luginaah, I.N.
Assessing the spatial distribution of nitrogen dioxide in London, Ontario, 1335

Ma, S.-C.
Experimental study on removals of SO_2 and NO_x using adsorption of activated carbon/microwave desorption, 1012

Ma, X.-Y.
Experimental study on removals of SO_2 and NO_x using adsorption of activated carbon/microwave desorption, 1012

Ma, Y.
An examination of boundary layer structure under the influence of the gap winds in Urumqi, China, during air pollution episode in winter, 26

MacDonnell, F.M.
Air impacts from three alternatives for producing JP-8 jet fuel, 1182

Analysis and comparison of inertinite-derived adsorbent with conventional adsorbents, 489

Maeda, H.
Transformation ofmicroflora during degradation of gaseous toluene in a biofilter detected using PCR-DGGE, 748

Maghirang, R.G.
Particulate matter emission rates from beef cattle feedlots in Kansas—Reverse dispersion modeling, 350

Mahmud, A.
Resolving the interactions between population density and air pollution emissions controls in the San Joaquin Valley, USA, 566

Majumdar, D.
Volatile organic compound emissions frommunicipal solidwaste disposal sites: Acase study of Mumbai, India, 398

Mancilla, Y.
A tunnel study to estimate emission factors from mobile sources in Monterrey, Mexico, 1431

Manomaiphiboon, K.
NO_2 and SO_2 dispersion modeling and relative roles of emission sources over Map Ta Phut industrial area, Thailand, 932

Ozone and its potential control strategy for Chon Buri city, Thailand, 1411

Mantilla Iglesias, E.
Characterization of the olfactory impact around a wastewater treatment plant: Optimization and validation of a hydrogen sulfide determination procedure based on passive diffusion sampling, 863

Estimation of hydrogen sulfide emission rates at several wastewater treatment plants through experimental concentration measurements and dispersion modeling, 758

Martinez, M.A.
Chemical characterization and factor analysis of $PM_{2.5}$ in two sites of Monterrey, Mexico, 817

McEwen, J.D.N.
Black carbon particulate matter emission factors for buoyancy-driven associated gas flares, 307

McGee, R.
Comparison of seasonal phenol and p-cresol emissions from ground-level area sources in a dairy operation in central Texas, 381

McKenna, J.
The evolution and impact of testing baghouse filter performance, 916

Mejia, G.M.
Chemical characterization and factor analysis of $PM_{2.5}$ in two sites of Monterrey, Mexico, 817

Mendoza, A.
A tunnel study to estimate emission factors from mobile sources in Monterrey, Mexico, 1431

Chemical characterization and factor analysis of $PM_{2.5}$ in two sites of Monterrey, Mexico, 817

Meneses, V.P.
A novel methodology for determining low-cost fine particulate matter street sweeping routes, 242

Mentese, S.
Multiple comparisons of organic, microbial, and fine particulate pollutants in typical indoor environments: Diurnal and seasonal variations, 1380

Miake-Lye, R.C.
Determination of the emissions from an aircraft auxiliary power unit (APU) during the Alternative Aviation Fuel Experiment (AAFEX), 420

Mihalic, J.N.
Assessment of heterogeneity of metal composition of fine particulate matter collected from eight U.S. counties using principal component analysis, 773

Miller, W.
Demonstration of fuel switching on oceangoing vessels in the Gulf of Mexico, 1093

Mobley, D.
Quantification of emission factor uncertainty, 287

Molenar, J.V.
Fine particulate matter and visibility in the Lake Tahoe Basin: Chemical characterization, trends, and source apportionment, 953

Morita, Y.
Optimizing nutrient supply in a rotatory-switching biofilter for toluene vapor treatment, 451

Transformation ofmicroflora during degradation of gaseous toluene in a biofilter detected using PCR-DGGE, 748

Mou, Z.
Effect of biocover equipped with a novel passive air diffusion system on microbial methane oxidation and community of methanotrophs, 278

Mukhtar, S.
Comparison of seasonal phenol and p-cresol emissions from ground-level area sources in a dairy operation in central Texas, 381

Mulholland, J.A.
Development of outcome-based, multipollutant mobile source indicators, 431

Muñoz, R.C.
Relative roles of emissions and meteorology in the diurnal pattern of urban PM_{10}: Analysis of the daylight saving time effect, 642

Murphy, J.P.
Particulate matter emission rates from beef cattle feedlots in Kansas—Reverse dispersion modeling, 350

Murtonen, T.
Study of Miller timing on exhaust emissions of a hydrotreated vegetable oil (HVO)-fueled diesel engine, 1305

Nakai, Y.
Middle-thermophilic sulfur-oxidizing bacteria *Thiomonas* sp. RAN5 strain for hydrogen sulfide removal, 38

Nakajima, J.
Optimizing nutrient supply in a rotatory-switching biofilter for toluene vapor treatment, 451

Nakao, S.
Aging of secondary organic aerosol from α-pinene ozonolysis: Roles of hydroxyl and nitrate radicals, 1359

Nenes, A.
Modeling an air pollution episode in northwestern United States: Identifying the effect of nitrogen oxide and volatile organic compound emission changes on air pollutants formation using direct sensitivity analysis, 1150

Ngo, K.Q.
Toluene vapor capture by activated carbon particles in a dual gas-solid cyclone system, 924

Ni, J.-Q.
Emissions monitoring at a deep-pit swine finishing facility: Research methods and system performance, 1264

Ni, M.-J.
Dioxins and their fingerprint in size-classified fly ash fractions from municipal solid waste incinerators in China—Mechanical grate and fluidized bed units, 717

Nishi, K.
Transformation ofmicroflora during degradation of gaseous toluene in a biofilter detected using PCR-DGGE, 748

Nolte, C.G.
Evaluating the effects of climate change on summertime ozone using a relative response factor approach for policymakers, 1061

Oda, T.
Integration of a nonmetallic electrostatic precipitator and a wet scrubber for improved removal of particles and corrosive gas cleaning in semiconductor manufacturing industries, 905

Oh, T.S.
A real-time monitoring and assessment method for calculation of total amounts of indoor air pollutants emitted in subway stations, 517

Oiamo, T.H.
Assessing the spatial distribution of nitrogen dioxide in London, Ontario, 1335

Okunishi, S.
Optimizing nutrient supply in a rotatory-switching biofilter for toluene vapor treatment, 451

Transformation ofmicroflora during degradation of gaseous toluene in a biofilter detected using PCR-DGGE, 748

Olaguer, E.P.
Near-source air quality impacts of large olefin flares, 978

The potential near-source ozone impacts of upstream oil and gas industry emissions, 966

Olatosi, A.
Assessment of a regulatory model's performance relative to large spatial heterogeneity in observed ozone in Houston, Texas, 696

Ollison, W.
Quantification of personal exposure concentrations to gasoline vehicle emissions in high-end exposure microenvironments: Effects of fuel and season, 1346

Omar, A.E.Kh.
Effects of foliar application with compost tea and filtrate biogas slurry liquid on yield and fruit quality of Washington navel orange *(Citrus sinenesis Osbeck)* trees, 767

Orešek, N.
Effect of thermal processing on the characteristics of incineration fly ash, 500

Pachon, J.E.
Development of outcome-based, multipollutant mobile source indicators, 431

Pala, M.
Temporal and spatial variations of polycyclic aromatic hydrocarbon concentrations around a coke oven plant, 1003

Pan, L.
Source attribution for mercury deposition in the contiguous United States: Regional difference and seasonal variation, 52

Pan, W.-P.
Mercury emission and plant uptake of trace elements during early stage of soil amendment using flue gas desulfurization materials, 139

Pan, Y.
Acid neutralization of precipitation in Northern China, 204

Park, B.H.
Influence of pleat geometry on filter cleaning in PTFE/glass composite filter, 1257

Park, J.-S.
Feasibility study on cross-linked biopolymeric concrete encapsulating selenium glass wastes, 898

Park, J.H.
A real-time monitoring and assessment method for calculation of total amounts of indoor air pollutants emitted in subway stations, 517

Park, K.H.
Effect of the Mn oxidation state and lattice oxygen in Mn-based TiO_2 catalysts on the low-temperature selective catalytic reduction of NO by NH_3, 1085

Park, Y.K.
Toluene vapor capture by activated carbon particles in a dual gas-solid cyclone system, 924

Parnell, Jr., C.B.
Comparison of seasonal phenol and p-cresol emissions from ground-level area sources in a dairy operation in central Texas, 381

Pattey, E.
Trends in primary particulate matter emissions from Canadian agriculture, 737

Pegues, A.H.
Efficacy of recent state implementation plans for 8-hour ozone, 252

Peng, R.D.
Assessment of heterogeneity of metal composition of fine particulate matter collected from eight U.S. counties using principal component analysis, 773

Pham, M.
The evolution and impact of testing baghouse filter performance, 916

Piccardo, M.T.
Temporal and spatial variations of polycyclic aromatic hydrocarbon concentrations around a coke oven plant, 1003

Pickett, D.A.
Effect of biogas generation on radon emissions from landfills receiving radium-bearing waste from shale gas development, 1040

Pollack, A.K.
Effects of improved spatial and temporal modeling of on-road vehicle emissions, 471

Pollock, M.C.
Speciated arsenic in air: Measurement methodology and risk assessment considerations, 2

Polyak, L.M.
Assessment of heterogeneity of metal composition of fine particulate matter collected from eight U.S. counties using principal component analysis, 773

Pongprueksa, P.
Source attribution for mercury deposition in the contiguous United States: Regional difference and seasonal variation, 52

Pouliot, G.
Quantification of emission factor uncertainty, 287

Prabamroong, T.
Ozone and its potential control strategy for Chon Buri city, Thailand, 1411

Prati, M.V.
Bioethanol–gasoline fuel blends: Exhaust emissions and morphological characterization of particulate from a moped engine, 888

Priest, J.W.
Air impacts from three alternatives for producing JP-8 jet fuel, 1182

Analysis and comparison of inertinite-derived adsorbent with conventional adsorbents, 489

Puccini, M.
Bioethanol–gasoline fuel blends: Exhaust emissions and morphological characterization of particulate from a moped engine, 888

Qi, L.
Aging of secondary organic aerosol from α-pinene ozonolysis: Roles of hydroxyl and nitrate radicals, 1359

Qiu, G.
Trends in primary particulate matter emissions from Canadian agriculture, 737

Qiu, Y.
Comparison of on-road emissions for B-0, B-10, and B-20 in transit buses, 443

Qu, Y.
Control strategies of atmospheric mercury emissions from coal-fired power plants in China, 576

Rad, A.Y.
Multiple comparisons of organic, microbial, and fine particulate pollutants in typical indoor environments: Diurnal and seasonal variation, 1380

Ramirez, D.
Analysis and comparison of inertinite-derived adsorbent with conventional adsorbents, 489

Ramos-Bonilla, J.P.
Assessment of heterogeneity of metal composition of fine particulate matter collected from eight U.S. counties using principal component analysis, 773

Rattigan, O.V.
Multiple-year black carbon measurements and source apportionment using Delta-C in Rochester, New York, 880

Ray, M.B.
Application of ultraviolet light-emitting diode photocatalysis to remove volatile organic compounds from indoor air, 1032

Razote, E.B.
Particulate matter emission rates from beef cattle feedlots in Kansas—Reverse dispersion modeling, 350

Reid, K.R.
Speciated arsenic in air: Measurement methodology and risk assessment considerations, 2

Reilly, E.O.
Optimization of a horizontal-flow biofilm reactor for the removal of methane at low temperatures, 1166

Ren, A.
Characteristics of penicillin bacterial residue, 485

Roberts, S.A.
Reducing ammonia emissions from laying-hen houses through dietary manipulation, 160

Rosati, J.A.
Thermal inactivation of Bacillus anthracis surrogate spores in a bench-scale enclosed landfill gas flare, 151

Roth, M.
Review of Singapore's air quality and greenhouse gas emissions: Current situation and opportunities, 625

Rule, A.M.
Assessment of heterogeneity of metal composition of fine particulate matter collected from eight U.S. counties using principal component analysis, 773

Russell, A.G.
Development of outcome-based, multipollutant mobile source indicators, 431

Development of risk-based air quality management strategies under impacts of climate change, 557

Modeling an air pollution episode in northwestern United States: Identifying the effect of nitrogen oxide and volatile organic compound emission changes on air pollutants formation using direct sensitivity analysis, 1150

Rynk, R.
Mitigation of odor causing emissions—Bench-scale investigation, 1423

Ščasný, M.
Cancer values of prevented fatalities (VPFs), one size does not fit all: The benefits of contaminated site cleanups in Italy, 783

Sagebiel, J.
Quantification of personal exposure concentrations to gasoline vehicle emissions in high-end exposure microenvironments: Effects of fuel and season, 1346

Saiyasitpanich, P.
NO_2 and SO_2 dispersion modeling and relative roles of emission sources over Map Ta Phut industrial area, Thailand, 932

Salathé, E.P.
Evaluating the effects of climate change on summertime ozone using a relative response factor approach for policymakers, 1061

Samec, N.
Effect of thermal processing on the characteristics of incineration fly ash, 500

SankaraRao, B.
A real-time monitoring and assessment method for calculation of total amounts of indoor air pollutants emitted in subway stations, 517

Sarjovaara, T.
Study of Miller timing on exhaust emissions of a hydrotreated vegetable oil (HVO)-fueled diesel engine, 1305

Sarnat, J.A.
Development of outcome-based, multipollutant mobile source indicators, 431

Sasivongpakdi, A.
Evaluation of a sustainable remediation option: Beneficial reuse of petroleum-contaminated sediment as an energy source, 1277

Sattler, M.L.
Air impacts from three alternatives for producing JP-8 jet fuel, 1182

Analysis and comparison of inertinite-derived adsorbent with conventional adsorbents, 489

Schauer, J. J.
Historical trends in the mass and chemical species concentrations of coarse particulate matter in the Los Angeles Basin and relation to sources and air quality regulations, 541

Schiappacasse, L.N.
Obtaining polycyclic aromatic hydrocarbon concentration ratios and molecular markers for residential wood combustion: Temuco, a case study, 44

Schuchmann, B.G.
Comparison of the MOVES2010a, MOBILE6.2, and EMFAC2007 mobile source emission models with on-road traffic tunnel and remote sensing measurements, 1134

Multispecies remote sensing measurements of vehicle emissions on ShermanWay in Van Nuys, California, 1127

Schultz, E.
A comparative study on the varying exposure to atmospheric fine and coarse particles under urban and rural conditions, 1313

Scungio, M.
Ultrafine particle emission from incinerators: The role of the fabric filter, 103

Seggiani, M.
Bioethanol–gasoline fuel blends: Exhaust emissions and morphological characterization of particulate from a moped engine, 888

Seila, R.L.
Near-road multipollutant profiles: Associations between volatile organic compounds and a tracer gas surrogate near a busy highway, 594

Shafer, M.M.
Historical trends in the mass and chemical species concentrations of coarse particulate matter in the Los Angeles Basin and relation to sources and air quality regulations, 541

Shang, G.
Effectiveness and mechanisms of hydrogen sulfide adsorption by camphor-derived biochar, 873

Shao, L.-M.
The emission patterns of volatile organic compounds during aerobic biotreatment of municipal solid waste using continuous and intermittent aeration, 461

Sharma, N.C.P.
An inexpensive active optical remote sensing instrument for assessing aerosol distributions, 198

Sharmin, R.
Application of ultraviolet light-emitting diode photocatalysis to remove volatile organic compounds from indoor air, 1032

Shaw, S.L.
Effects of improved spatial and temporal modeling of on-road vehicle emissions, 471

Shen, G.
Effectiveness and mechanisms of hydrogen sulfide adsorption by camphor-derived biochar, 873

Shen, Z.-X.
Winter and Summer $PM_{2.5}$ Chemical Compositions in Fourteen Chinese Cities, 1214

Shetty, K.V.
A real-time monitoring and assessment method for calculation of total amounts of indoor air pollutants emitted in subway stations, 517

Shetty, S.K.
Source attribution for mercury deposition in the contiguous United States: Regional difference and seasonal variation, 52

Shih, H.-Y.
Development of regenerative dye impregnated mesoporous silica materials for assessing exposure to ammonia, 838

Shimaoka, T.
Atmospheric mercury emissions from waste combustions measured by continuous monitoring devices, 686

Shores, R.C.
Near-road multipollutant profiles: Associations between volatile organic compounds and a tracer gas surrogate near a busy highway, 594

Sigaev, V.I.
Awind tunnel test of newly developed personal bioaerosol samplers, 828

Sioutas, C.
Historical trends in the mass and chemical species concentrations of coarse particulate matter in the Los Angeles Basin and relation to sources and air quality regulations, 541

Sistani, K. R.
Mercury emission and plant uptake of trace elements during early stage of soil amendment using flue gas desulfurization materials, 139

Sridhar, M.K.C.
Polycyclic aromatic hydrocarbon burden in ambient air in selected Niger Delta communities in Nigeria, 18

Srivastava, A.
Volatile organic compound emissions frommunicipal solidwaste disposal sites: Acase study of Mumbai, India, 398

Stabile, L.
Ultrafine particle emission from incinerators: The role of the fabric filter, 103

Stedman, D.H.
Comparison of the MOVES2010a, MOBILE6.2, and EMFAC2007 mobile source emission models with on-road traffic tunnel and remote sensing measurements, 1134

Multispecies remote sensing measurements of vehicle emissions on ShermanWay in Van Nuys, California, 1127

Stella, A.
Temporal and spatial variations of polycyclic aromatic hydrocarbon concentrations around a coke oven plant, 1003

Su, M.-C.
Potential assessment of establishing a renewable energy plant in a rural agricultural area, 662

Su, W.-C.
A wind tunnel test of newly developed personal bioaerosol samplers, 828

Sukhapan, J.
Ozone and its potential control strategy for Chon Buri city, Thailand, 1411

Sun, Z.
Effects of activated carbon surface properties on the adsorption of volatile organic compounds, 1196

Swan, N.
A new approach to characterize emission contributions from area sources during optical remote sensing technique testing, 1403

Methane emissions from 20 landfills across the United States using vertical radial plume mapping, 183

Switzer, D.
Ambient measurements of 2,2,4-trimethyl, 1,3-pentanediol monoisobutyrate in Southern California, 680

Tacina, K.B.
Determination of the emissions from an aircraft auxiliary power unit (APU) during the Alternative Aviation Fuel Experiment (AAFEX), 420

Tagaris, E.
Development of risk-based air quality management strategies under impacts of climate change, 557

Takahashi, F.
Atmospheric mercury emissions from waste combustions measured by continuous monitoring devices, 686

Tamura, K.
Driver exposure to particulate matter in Bangkok, 64

Tang, J.-F.
The emission patterns of volatile organic compounds during aerobic biotreatment of municipal solid waste using continuous and intermittent aeration, 461

Tang, K.
Assessing the spatial distribution of nitrogen dioxide in London, Ontario, 1335

Tang, S.
The addition of modified attapulgite reduces the emission of nitrous oxide and ammonia from aerobically composted chicken manure, 1174

Thepanondh, S.
NO_2 and SO_2 dispersion modeling and relative roles of emission sources over Map Ta Phut industrial area, Thailand, 932

Thoma, E.D.
Near-road multipollutant profiles: Associations between volatile organic compounds and a tracer gas surrogate near a busy highway, 594

Thornhill, K.L.
Determination of the emissions from an aircraft auxiliary power unit (APU) during the Alternative Aviation Fuel Experiment (AAFEX), 420

Tian, H.
Control strategies of atmospheric mercury emissions from coal-fired power plants in China, 576

Tian, Y.
Rapidly eliminating pathogenic microorganisms in large air space using spraying •OH radicals, 393

Tie, X.-X.
Winter and Summer $PM_{2.5}$ Chemical Compositions in Fourteen Chinese Cities, 1214

Timko, M.T.
Determination of the emissions from an aircraft auxiliary power unit (APU) during the Alternative Aviation Fuel Experiment (AAFEX), 420

Tirler, W.
Ultrafine particle emission from incinerators: The role of the fabric filter, 103

Tolbert, P.E.
Development of outcome-based, multipollutant mobile source indicators, 431

Tolchinsky, A.D.
Awind tunnel test of newly developed personal bioaerosol samplers, 828

Tong, H.
The effects of manganese precursors on Mn-based/TiO_2 catalysts for catalytic reduction of NO with NH_3, 271

Tonin, S.
Cancer values of prevented fatalities (VPFs), one size does not fit all: The benefits of contaminated site cleanups in Italy, 783

Trail, M.
Modeling an air pollution episode in northwestern United States: Identifying the effect of nitrogen oxide and volatile organic compound emission changes on air pollutants formation using direct sensitivity analysis, 1150

Tsai, H.-H.
Enhanced mercuric chloride adsorption onto sulfur-modified activated carbons derived from waste tires, 799

Tsai, J.-H.
Experimental studies of the dilution of vehicle exhaust pollutants by environment-protecting pervious pavement, 92

Tsimpidi, A.P.
Modeling an air pollution episode in northwestern United States: Identifying the effect of nitrogen oxide and volatile organic compound emission changes on air pollutants formation using direct sensitivity analysis, 1150

Tufts, J.A.M.
Thermal inactivation of Bacillus anthracis surrogate spores in a bench-scale enclosed landfill gas flare, 151

Utell, M.J.
Multiple-year black carbon measurements and source apportionment using Delta-C in Rochester, New York, 880

Valerio, F.
Temporal and spatial variations of polycyclic aromatic hydrocarbon concentrations around a coke oven plant, 1003

Valesco, E.
Review of Singapore's air quality and greenhouse gas emissions: Current situation and opportunities, 625

Veawab, A.
A generalized fuzzy linear programming approach for environmental management problem under uncertainty, 72

Vidal, V.
Obtaining polycyclic aromatic hydrocarbon concentration ratios and molecular markers for residential wood combustion: Temuco, a case study, 44

Virtanen, A.
Study of Miller timing on exhaust emissions of a hydrotreated vegetable oil (HVO)-fueled diesel engine, 1305

Vitolo, S.
Bioethanol–gasoline fuel blends: Exhaust emissions and morphological characterization of particulate from a moped engine, 888

Vizuete, W.
Assessment of a regulatory model's performance relative to large spatial heterogeneity in observed ozone in Houston, Texas, 696

Walsh, R.
Optimization of a horizontal-flow biofilm reactor for the removal of methane at low temperatures, 1166

Walter, G.R.
Effect of biogas generation on radon emissions from landfills receiving radium-bearing waste from shale gas development, 1040

Wang, G.
Enhanced mercuric chloride adsorption onto sulfur-modified activated carbons derived from waste tires, 799

Wang, G.-H.
Winter and Summer $PM_{2.5}$ Chemical Compositions in Fourteen Chinese Cities, 1214

Wang, H.
Effect of biocover equipped with a novel passive air diffusion system on microbial methane oxidation and community of methanotrophs, 278

Preparation, characterization, and phenol adsorption of activated carbons from oxytetracycline bacterial residue, 1394

Wang, J.
Adsorption Behavior of Toluene on Modified 1X Molecular Sieves, 1227

Wang, L.
Comparison of the purification performance and microbial community functional diversity in flowdirectional- switching and unidirectional-flow biotrickling filters, 1203

Wang, T.
Effectiveness and mechanisms of hydrogen sulfide adsorption by camphor-derived biochar, 873

Wang, X.
Comparison of six particle size distribution models on the goodness-of-fit to particulate matter sampled from animal buildings, 725

Wang, Y.
Acid neutralization of precipitation in Northern China, 204

Control strategies of atmospheric mercury emissions from coal-fired power plants in China, 576

Multiple-year black carbon measurements and source apportionment using Delta-C in Rochester, New York, 880

Wang, Y.-W.
Mercury emission and plant uptake of trace elements during early stage of soil amendment using flue gas desulfurization materials, 139

Wang-Li, L.
Field evaluation of particulate matter measurements using tapered element oscillating microbalance in a layer house, 322

Watson, J.G.
Energy supplies and future engines for land, sea, and air, 1233

Winter and Summer $PM_{2.5}$ Chemical Compositions in Fourteen Chinese Cities, 1214

Weedon, J.
Ambient air pollution and term birth weight in Texas from 1998 to 2004, 1285

Wey, C.
Determination of the emissions from an aircraft auxiliary power unit (APU) during the Alternative Aviation Fuel Experiment (AAFEX), 420

Whitefield, P.
Determination of the emissions from an aircraft auxiliary power unit (APU) during the Alternative Aviation Fuel Experiment (AAFEX), 420

Wiedinmyer, C.
Evaluating the effects of climate change on summertime ozone using a relative response factor approach for policymakers, 1061

Wilson, D.G.
Energy supplies and future engines for land, sea and air, 607

Wilson, R.
Efficacy of recent state implementation plans for 8-hour ozone, 252

Winstead, E.
Determination of the emissions from an aircraft auxiliary power unit (APU) during the Alternative Aviation Fuel Experiment (AAFEX), 420

Wisner, E.
Quantification of emission factor uncertainty, 287

Wolf-Benning, U.
A comparative study on the varying exposure to atmospheric fine and coarse particles under urban and rural conditions, 1313

Wong, D.C.
Sensitivity and linearity analysis of ozone in East Asia: The effects of domestic emission and intercontinental transport, 1102

Wood, E.C.
Determination of the emissions from an aircraft auxiliary power unit (APU) during the Alternative Aviation Fuel Experiment (AAFEX), 420

Wu, C.-F.
Evaluating the performance of the horizontal radial plume mapping technique for locating multiple plumes, 1249

Wu, D.
Acid neutralization of precipitation in Northern China, 204

Xia, X.
An examination of boundary layer structure under the influence of the gap winds in Urumqi, China, during air pollution episode in winter, 26

Xie, K.
The addition of modified attapulgite reduces the emission of nitrous oxide and ammonia from aerobically composted chicken manure, 1174

Xin, H.
Reducing ammonia emissions from laying-hen houses through dietary manipulation, 160

Xin, Y.
An examination of boundary layer structure under the influence of the gap winds in Urumqi, China, during air pollution episode in winter, 26

Xu, J.
Comparison of the purification performance and microbial community functional diversity in flowdirectional- switching and unidirectional-flow biotrickling filters, 1203

Determination of partition and diffusion coefficients of formaldehyde in selected building materials and impact of relative humidity, 671

Xu, P.
The addition of modified attapulgite reduces the emission of nitrous oxide and ammonia from aerobically composted chicken manure, 1174

Xu, X.
Assessing the spatial distribution of nitrogen dioxide in London, Ontario, 1335

Xue, S.-H.
Enhanced mercuric chloride adsorption onto sulfur-modified activated carbons derived from waste tires, 799

Xue, Z.
Control strategies of atmospheric mercury emissions from coal-fired power plants in China, 576

Yan, F.
Removal of nitric oxide in a biotrickling filter under thermophilic condition using *Chelatococcus daeguensis*, 509

Yan, J.-H.
Dioxins and their fingerprint in size-classified fly ash fractions from municipal solid waste incinerators in China—Mechanical grate and fluidized bed units, 717

Yang, J.
An examination of boundary layer structure under the influence of the gap winds in Urumqi, China, during air pollution episode in winter, 26

Yang, N.
The emission patterns of volatile organic compounds during aerobic biotreatment of municipal solid waste using continuous and intermittent aeration, 461

Yang, S.
The addition of modified attapulgite reduces the emission of nitrous oxide and ammonia from aerobically composted chicken manure, 1174

Yang, X.
An examination of boundary layer structure under the influence of the gap winds in Urumqi, China, during air pollution episode in winter, 26

Comparison of six particle size distribution models on the goodness-of-fit to particulate matter sampled from animal buildings, 725

Yang, Z.
Application of neutral electrolyzed water spray for reducing dust levels in a layer breeding house, 1329

Yano, E.
Driver exposure to particulate matter in Bangkok, 64

Yao, J.-J.
Experimental study on removals of SO_2 and NO_x using adsorption of activated carbon/microwave desorption, 1012

Yeardley, Jr., R.B.
Evaluation of a sustainable remediation option: Beneficial reuse of petroleum-contaminated sediment as an energy source, 1277

Yen, M.-T.
Experimental studies of the dilution of vehicle exhaust pollutants by environment-protecting pervious pavement, 92

Yen, T.F.
Feasibility study on cross-linked biopolymeric concrete encapsulating selenium glass wastes, 898

Yin, N.
Comparison of the purification performance and microbial community functional diversity in flowdirectional- switching and unidirectional-flow biotrickling filters, 1203

Yoa, S.-J.
Integration of a nonmetallic electrostatic precipitator and a wet scrubber for improved removal of particles and corrosive gas cleaning in semiconductor manufacturing industries, 905

Yoo, C.K.
A real-time monitoring and assessment method for calculation of total amounts of indoor air pollutants emitted in subway stations, 517

Yu, J.Z.
Carbonyl emissions from vehicular exhausts sources in Hong Kong, 221

Yu, K.-P.
Enhancement of the deposition of ultrafine secondary organic aerosols by the negative air ion and the effect of relative humidity, 1296

Yu, W.
Acid neutralization of precipitation in Northern China, 204

Yu, Y.
Adsorption Behavior of Toluene on Modified 1X Molecular Sieves, 1227

Yu, Y.-P.
Breakthrough indicator for aromatic VOCs using needle trap samplers for activated carbon adsorbent, 946

Yu, Z.
Determination of the emissions from an aircraft auxiliary power unit (APU) during the Alternative Aviation Fuel Experiment (AAFEX), 420

Yuan, C.-S.
Enhanced mercuric chloride adsorption onto sulfur-modified activated carbons derived from waste tires, 799

Yucel, A.
Mitigation of odor causing emissions—Bench-scale investigation, 1423

Yusoff, S.
Benefits of clean development mechanism application on the life cycle assessment perspective: A case study in the palm oil industry, 299

Zhang, F.
The addition of modified attapulgite reduces the emission of nitrous oxide and ammonia from aerobically composted chicken manure, 1174

Zhang, G.
Application of neutral electrolyzed water spray for reducing dust levels in a layer breeding house, 1329

Zhang, J.S.
Determination of partition and diffusion coefficients of formaldehyde in selected building materials and impact of relative humidity, 671

Zhang, R.
Removal of nitric oxide in a biotrickling filter under thermophilic condition using *Chelatococcus daeguensis*, 509

Zhang, Y.
Comparison of six particle size distribution models on the goodness-of-fit to particulate matter sampled from animal buildings, 725

Evaluating the effects of climate change on summertime ozone using a relative response factor approach for policymakers, 1061

Zhang, Z.
Rapidly eliminating pathogenic microorganisms in large air space using spraying •OH radicals, 393

Zhao, W.
Characteristics of penicillin bacterial residue, 485

Zhao, Y.
Experimental study on removals of SO_2 and NO_x using adsorption of activated carbon/microwave desorption, 1012

Zheng, L.
Adsorption Behavior of Toluene on Modified 1X Molecular Sieves, 1227

Zheng, W.
Application of neutral electrolyzed water spray for reducing dust levels in a layer breeding house, 1329

Zhou, B.
Preparation, characterization, and phenol adsorption of activated carbons from oxytetracycline bacterial residue, 1394

Zhu, N.
Preparation, characterization, and phenol adsorption of activated carbons from oxytetracycline bacterial residue, 1394

Zielinska, B.
Comparison of the MOVES2010a, MOBILE6.2, and EMFAC2007 mobile source emission models with on-road traffic tunnel and remote sensing measurements, 1134

Quantification of personal exposure concentrations to gasoline vehicle emissions in high-end exposure microenvironments: Effects of fuel and season, 1346

Zupanič, Franc
Effect of thermal processing on the characteristics of incineration fly ash, 500